

Anti-HEAB Rabbit Monoclonal Antibody

Catalog # ABO16168

Product Information

Application	WB, IHC, FC
Primary Accession	Q92989
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-HEAB Rabbit Monoclonal Antibody . Tested in WB, IHC, Flow Cytometry applications. This antibody reacts with Human.


Additional Information

Gene ID	10978
Other Names	Polyribonucleotide 5'-hydroxyl-kinase Clp1 {ECO:0000255 HAMAP-Rule:MF_03035}, 2.7.1.78 {ECO:0000255 HAMAP-Rule:MF_03035}, Polyadenylation factor Clp1 {ECO:0000255 HAMAP-Rule:MF_03035}, Polynucleotide kinase Clp1 {ECO:0000255 HAMAP-Rule:MF_03035}, Pre-mRNA cleavage complex II protein Clp1 {ECO:0000255 HAMAP-Rule:MF_03035}, CLP1 {ECO:0000255 HAMAP-Rule:MF_03035}, HEAB
Calculated MW	47646
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 27C47
Immunogen	A synthesized peptide derived from human HEAB
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CLP1 {ECO:0000255 HAMAP-Rule:MF_03035}
Synonyms	HEAB
Function	Polynucleotide kinase that can phosphorylate the 5'-hydroxyl groups of double-stranded RNA (dsRNA), single-stranded RNA (ssRNA), double-stranded DNA (dsDNA) and double-stranded DNA:RNA hybrids. dsRNA is phosphorylated more efficiently than dsDNA, and the RNA component of a DNA:RNA hybrid is phosphorylated more efficiently than the DNA component. Plays a key role in both tRNA splicing and mRNA 3'-end formation. Component of the tRNA splicing endonuclease complex: phosphorylates the 5'-terminus of the tRNA 3'-exon during tRNA splicing; this phosphorylation event is a prerequisite for the subsequent ligation of the two exon halves and the production of a mature tRNA (PubMed:24766809, PubMed:24766810). Its role in tRNA splicing and maturation is required for cerebellar development (PubMed:24766809, PubMed:24766810). Component of the pre-mRNA cleavage complex II (CF-II), which seems to be required for mRNA 3'-end formation. Also phosphorylates the 5'-terminus of exogenously introduced short interfering RNAs (siRNAs), which is a necessary prerequisite for their incorporation into the RNA-induced silencing complex (RISC). However, endogenous siRNAs and microRNAs (miRNAs) that are produced by the cleavage of dsRNA precursors by DICER1 already contain a 5'-phosphate group, so this protein may be dispensible for normal RNA-mediated gene silencing.
Cellular Location	Nucleus {ECO:0000255 HAMAP-Rule:MF_03035, ECO:0000269 PubMed:11060040, ECO:0000269 PubMed:24766810}

Images



Western blot analysis of HEAB expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.