

Anti-C1s Rabbit Monoclonal Antibody

Catalog # ABO16197

Product Information

Application	WB, IHC, IP
Primary Accession	P09871
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-C1s Rabbit Monoclonal Antibody . Tested in WB, IHC, IP applications. This antibody reacts with Human.

Additional Information

Gene ID	716
Other Names	Complement C1s subcomponent, 3.4.21.42, C1 esterase, Complement component 1 subcomponent s, Complement C1s subcomponent heavy chain, Complement C1s subcomponent light chain, C1S
Calculated MW	76684
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 27C76 A synthesized peptide derived from human C1s
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	C1S {ECO:0000303 PubMed:3500856, ECO:0000312 HGNC:HGNC:1247}
Function	Component of the complement C1 complex, a multiprotein complex that initiates the classical pathway of the complement system, a cascade of

proteins that leads to phagocytosis and breakdown of pathogens and signaling that strengthens the adaptive immune system (PubMed:[11445589](#), PubMed:[16169853](#), PubMed:[417728](#), PubMed:[467643](#), PubMed:[6271784](#), PubMed:[6282646](#), PubMed:[6319179](#), PubMed:[70787](#), PubMed:[9422791](#)). C1S is activated following association of the C1 complex with immunoglobulins (IgG or IgM) complexed with antigens to form antigen-antibody complexes on the surface of pathogens (PubMed:[34155115](#)). C1S is cleaved and activated by C1R to generate C1s subcomponent heavy and light chains (PubMed:[11445589](#), PubMed:[6271784](#)). C1s subcomponent light chain then cleaves and activates C2 and C4, the next components of the classical complement pathway (PubMed:[16169853](#), PubMed:[467643](#), PubMed:[6282646](#), PubMed:[6319179](#), PubMed:[6906228](#), PubMed:[70787](#), PubMed:[9422791](#)).

Cellular Location

Secreted. Cell surface. Note=Recruited to the surface of pathogens by the C1Q subcomplex.

Images



Western blot analysis of C1s expression in human plasma cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.