

Anti-Sterol carrier protein 2 Rabbit Monoclonal Antibody

Catalog # ABO16226

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P22307
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Sterol carrier protein 2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	6342
Other Names	Sterol carrier protein 2, SCP-2, Acetyl-CoA C-myristoyltransferase, 2.3.1.155, SCP-chi, SCPX, Sterol carrier protein X, SCP-X, SCP2 (HGNC:10606)
Calculated MW	58994
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 28S05
Immunogen	A synthesized peptide derived from human Sterol carrier protein 2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	SCP2 (HGNC:10606)
Function	[Isoform SCPx]: Plays a crucial role in the peroxisomal oxidation of branched-chain fatty acids (PubMed: 10706581). Catalyzes the last step of the peroxisomal beta-oxidation of branched chain fatty acids and the side chain

of the bile acid intermediates di- and trihydroxycoprostanic acids (DHCA and THCA) (PubMed:[10706581](#)). Also active with medium and long straight chain 3-oxoacyl-CoAs. Stimulates the microsomal conversion of 7-dehydrocholesterol to cholesterol and transfers phosphatidylcholine and 7-dehydrocholesterol between membranes, in vitro (By similarity). Isoforms SCP2 and SCPx cooperate in peroxisomal oxidation of certain naturally occurring tetramethyl- branched fatty acyl-CoAs (By similarity).

Cellular Location

[Isoform SCP2]: Peroxisome {ECO:0000250|UniProtKB:P32020}. Cytoplasm. Mitochondrion. Endoplasmic reticulum {ECO:0000250|UniProtKB:P32020}. Mitochondrion {ECO:0000250|UniProtKB:P32020}

Tissue Location

Liver, fibroblasts, and placenta.

Images




Figure 1. Western blot analysis of Sterol carrier protein 2 using anti-Sterol carrier protein 2 antibody (M02947-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,

Lane 2: human HUH-7 whole cell lysates,

Lane 3: rat RH35 whole cell lysates,

Lane 4: mouse liver tissue lysates,

Lane 5: mouse HEPA1-6 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Sterol carrier protein 2 antigen affinity purified monoclonal antibody (Catalog # M02947-1) at 1:1000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Sterol carrier protein 2 at approximately 59 kDa. The expected band size for Sterol carrier protein 2 is at 59 kDa.




Figure 2. IF analysis of Sterol carrier protein 2 using anti-Sterol carrier protein 2 antibody (M02947-1). Sterol carrier protein 2 was detected in an immunocytochemical section of Hela cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated at 1:50 rabbit anti-Sterol carrier protein 2 Antibody (M02947-1) overnight at 4°C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.