

Anti-Phospho-Rad17 (S656) Rabbit Monoclonal Antibody

Catalog # ABO16294

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	O75943
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Phospho-Rad17 (S656) Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

Additional Information

Gene ID	5884
Other Names	Cell cycle checkpoint protein RAD17, hRad17, RF-C/activator 1 homolog, RAD17 {ECO:0000303 PubMed:9878245, ECO:0000312 HGNC:HGNC:9807}
Calculated MW	77055
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 28R30
Immunogen	A synthesized peptide derived from human Phospho-Rad17 (S656)
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	RAD17 {ECO:0000303 PubMed:9878245, ECO:0000312 HGNC:HGNC:9807}
Function	Essential for sustained cell growth, maintenance of chromosomal stability, and ATR-dependent checkpoint activation upon DNA damage (PubMed: 10208430 , PubMed: 11418864 , PubMed: 11687627 ,

PubMed:[11799063](#), PubMed:[12672690](#), PubMed:[14624239](#), PubMed:[15235112](#)). Has a weak ATPase activity required for binding to chromatin (PubMed:[10208430](#), PubMed:[11418864](#), PubMed:[11687627](#), PubMed:[11799063](#), PubMed:[12672690](#), PubMed:[14624239](#), PubMed:[15235112](#)). Participates in the recruitment of the 9-1-1 (RAD1-RAD9-HUS1) complex and RHNO1 onto chromatin, and in CHEK1 activation (PubMed:[21659603](#)). Involved in homologous recombination by mediating recruitment of the MRN complex to DNA damage sites (PubMed:[24534091](#)). May also serve as a sensor of DNA replication progression (PubMed:[12578958](#), PubMed:[14500819](#), PubMed:[15538388](#)).


Cellular Location

Nucleus. Chromosome Note=Phosphorylated form redistributes to discrete nuclear foci upon DNA damage (PubMed:11799063). Localizes to DNA double-strand breaks (DSBs) (PubMed:24534091).

Tissue Location

Overexpressed in various cancer cell lines and in colon carcinoma (at protein level). Isoform 2 and isoform 3 are the most abundant isoforms in non irradiated cells (at protein level) Ubiquitous at low levels. Highly expressed in testis, where it is expressed within the germinal epithelium of the seminiferous tubuli Weakly expressed in seminomas (testicular tumors)

Images



Western blot analysis of Phospho-Rad17 (S656) expression in (1) HeLa cell treated with CA lysate; (2) HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.