

Anti-DENN Rabbit Monoclonal Antibody

Catalog # ABO16350

Product Information

Application	WB, IHC
Primary Accession	Q8WVG6
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-DENN Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	8567
Other Names	MAP kinase-activating death domain protein, Differentially expressed in normal and neoplastic cells, Insulinoma glucagonoma clone 20, Rab3 GDP/GTP exchange factor, RabGEF, Rab3 GDP/GTP exchange protein, Rab3GEP, MADD {ECO:0000312 EMBL:AAB57735.1, ECO:0000312 HGNC:HGNC:6766}
Calculated MW	183303
Application Details	WB 1:500-1:2000 IHC 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 28M86
Immunogen	A synthesized peptide derived from human DENN
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	MADD {ECO:0000312 EMBL:AAB57735.1, ECO:0000312 HGNC:HGNC:6766}
Function	Guanyl-nucleotide exchange factor that regulates small GTPases of the Rab

family (PubMed:[18559336](#), PubMed:[20937701](#)). Converts GDP-bound inactive form of RAB27A and RAB27B to the GTP-bound active forms (PubMed:[18559336](#), PubMed:[20937701](#)). Converts GDP-bound inactive form of RAB3A, RAB3C and RAB3D to the GTP-bound active forms, GTPases involved in synaptic vesicle exocytosis and vesicle secretion (By similarity). Plays a role in synaptic vesicle formation and in vesicle trafficking at the neuromuscular junction (By similarity). Involved in up-regulating a post-docking step of synaptic exocytosis in central synapses (By similarity). Probably by binding to the motor proteins KIF1B and KIF1A, mediates motor-dependent transport of GTP-RAB3A- positive vesicles to the presynaptic nerve terminals (By similarity). Plays a role in TNFA-mediated activation of the MAPK pathway, including ERK1/2 (PubMed:[32761064](#)). May link TNFRSF1A with MAP kinase activation (PubMed:[9115275](#)). May be involved in the regulation of TNFA-induced apoptosis (PubMed:[11577081](#), PubMed:[32761064](#)).


Cellular Location

Cell membrane. Cytoplasm. Cell projection, axon
{ECO:0000250|UniProtKB:Q80U28}

Tissue Location

Expressed in testis, ovary, brain and heart (PubMed:8988362). Expressed in spleen, thymus, prostate, testis, ovary, small intestine and colon (PubMed:9115275). Expressed in liver (PubMed:9796103). [Isoform 2]: Expressed in the brain, breast, kidney, lung, ovary, pancreas, testis, uterus, stomach and thyroid [Isoform 4]: Expressed in the brain, breast, kidney, lung, ovary, pancreas, testis, uterus, stomach and thyroid [Isoform 6]: Not detected in the brain, breast, kidney, lung, ovary, pancreas, testis, uterus, stomach and thyroid

Images



Western blot analysis of DENN expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.