10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CTP synthase / CTPS Rabbit Monoclonal Antibody

Catalog # ABO16373

Product Information

Application	WB, FC
Primary Accession	P17812
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CTP synthase / CTPS Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse.

Additional Information

Gene ID	1503
Other Names	CTP synthase 1, 6.3.4.2, CTP synthetase 1, UTP--ammonia ligase 1, CTPS1 (HGNC:2519)
Calculated MW	66690
Application Details	WB 1:500-1:2000 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 29C09
Immunogen	A synthesized peptide derived from human CTP synthase / CTPS
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CTPS1 {ECO:0000303|PubMed:16179339, ECO:0000312|HGNC:HGNC:2519}
Function	CTP synthase involved in the de novo synthesis of CTP, a precursor of DNA, RNA and phospholipids (PubMed:16179339, PubMed:17189248, PubMed:17463002, PubMed:24870241, PubMed:28459447, PubMed:34583994). Catalyzes the ATP-dependent amination of UTP to CTP

with either L-glutamine or ammonia as a source of nitrogen (PubMed:16179339, PubMed:24870241, PubMed:28459447, PubMed:34583994). CTPS1 CTP synthase activity plays a crucial role in the proliferation of activated lymphocytes and immunity; additional CTP being required to meet increased demand for DNA, RNA and lipid membrane biosynthesis in proliferating lymphocytes (PubMed:24870241, PubMed:8530356). In addition to CTP synthase activity, also acts as a protein deamidase that catalyzes the side chain deamidation of specific asparagine residues of proteins to aspartate (PubMed:40240600). Acts as a negative regulator of innate immunity by mediating deamidation of 'Asn-85' of IRF3, preventing IRF3 from binding DNA (By similarity). Facilitates chromatin relaxation in response to DNA damage by mediating deamidation of 'Asn-76' and 'Asn-77' of histone H1, thereby promoting subsequent acetylation of histone H1 at 'Lys-75' (H1K75ac), increasing chromatin accessibility to facilitate the recruitment of DNA repair proteins (PubMed:40240600).


Cellular Location

Cytoplasm, cytosol. Nucleus. Chromosome Note=Cytoplasmic and nuclear; when active, detected in long filamentous structures named cytoophidium (PubMed:24503052, PubMed:25223282, PubMed:28459447). Enriched on chromatin in response to DNA damage (PubMed:40240600).

Tissue Location

Widely expressed at low level, except in activated T-cells where it is highly expressed.

Images



Western blot analysis of CTP synthase / CTPS expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.