10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-PLSCR3 Rabbit Monoclonal Antibody

Catalog # ABO16377

Product Information

Application	WB
Primary Accession	Q9NRY6
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-PLSCR3 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	57048
Other Names	Phospholipid scramblase 3, PL scramblase 3, Ca(2+)-dependent phospholipid scramblase 3, PLSCR3
Calculated MW	31648
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 29P13
Immunogen	A synthesized peptide derived from human PLSCR3
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	PLSCR3
Function	Catalyzes calcium-induced ATP-independent rapid bidirectional and non-specific movement of the phospholipids (lipid scrambling or lipid flip-flop) between the inner and outer membrane of the mitochondria (PubMed:14573790, PubMed:17226776, PubMed:18358005, PubMed:29337693, PubMed:31769662). Plays an important role in

mitochondrial respiratory function, morphology, and apoptotic response (PubMed:12649167, PubMed:14573790, PubMed:17226776, PubMed:18358005). Mediates the translocation of cardiolipin from the mitochondrial inner membrane to outer membrane enhancing t-Bid induced cytochrome c release and apoptosis (PubMed:14573790, PubMed:17226776, PubMed:18358005). Enhances TNFSF10-induced apoptosis by regulating the distribution of cardiolipin in the mitochondrial membrane resulting in increased release of apoptogenic factors and consequent amplification of the activity of caspases (PubMed:18491232). Regulates cardiolipin de novo biosynthesis and its resynthesis (PubMed:16939411).


Cellular Location

Mitochondrion membrane; Single-pass type II membrane protein {ECO:0000250|UniProtKB:Q6QBQ4}. Mitochondrion inner membrane {ECO:0000250|UniProtKB:Q6QBQ4}; Single-pass type II membrane protein {ECO:0000250|UniProtKB:Q6QBQ4}. Nucleus {ECO:0000250|UniProtKB:Q9JIZ9} Note=Palmitoylation regulates its localization to the cell membrane or the nucleus; trafficking to the cell membrane is dependent upon palmitoylation whereas in the absence of palmitoylation, localizes to the nucleus. {ECO:0000250|UniProtKB:Q9JIZ9}

Tissue Location

Expressed in heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, uterus, small intestine and peripheral blood lymphocytes. Not detected in testis, brain and liver

Images



Western blot analysis of PLSCR3 expression in BxPC 3 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.