

Anti-ITPK1 Rabbit Monoclonal Antibody

Catalog # ABO16414

Product Information

Application	WB, IHC
Primary Accession	Q13572
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-ITPK1 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

Additional Information

Gene ID	3705
Other Names	Inositol-tetrakisphosphate 1-kinase, 2.7.1.134, Inositol 1, 3, 4-trisphosphate 5/6-kinase, Inositol-triphosphate 5/6-kinase, Ins(1, 3, 4)P(3) 5/6-kinase, 2.7.1.159, ITPK1 (HGNC:6177)
Calculated MW	45621
Application Details	WB 1:500-1:2000 IHC 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 29I50 A synthesized peptide derived from human ITPK1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	ITPK1 (HGNC:6177)
Function	Kinase that can phosphorylate various inositol polyphosphate such as Ins(3,4,5,6)P4 or Ins(1,3,4)P3 (PubMed: 11042108 , PubMed: 8662638). Phosphorylates Ins(3,4,5,6)P4 at position 1 to form Ins(1,3,4,5,6)P5

(PubMed:[11042108](#)). This reaction is thought to have regulatory importance, since Ins(3,4,5,6)P4 is an inhibitor of plasma membrane Ca(2+)-activated Cl(-) channels, while Ins(1,3,4,5,6)P5 is not. Also phosphorylates Ins(1,3,4)P3 on O-5 and O-6 to form Ins(1,3,4,6)P4, an essential molecule in the hexakisphosphate (InsP6) pathway (PubMed:[11042108](#), PubMed:[8662638](#)). Also acts as an inositol polyphosphate phosphatase that dephosphorylates Ins(1,3,4,5)P4 and Ins(1,3,4,6)P4 to Ins(1,3,4)P3, and Ins(1,3,4,5,6)P5 to Ins(3,4,5,6)P4 (PubMed:[11909533](#), PubMed:[17616525](#)). May also act as an isomerase that interconverts the inositol tetrakisphosphate isomers Ins(1,3,4,5)P4 and Ins(1,3,4,6)P4 in the presence of ADP and magnesium (PubMed:[11909533](#)). Probably acts as the rate-limiting enzyme of the InsP6 pathway. Modifies TNF-induced apoptosis by interfering with the activation of TNFRSF1A-associated death domain (PubMed:[11909533](#), PubMed:[12925536](#), PubMed:[17616525](#)). Plays an important role in MLKL-mediated necroptosis. Produces highly phosphorylated inositol phosphates such as inositolhexakisphosphate (InsP6) which bind to MLKL mediating the release of an N-terminal auto-inhibitory region leading to its activation. Essential for activated phospho-MLKL to oligomerize and localize to the cell membrane during necroptosis (PubMed:[17616525](#)).

Tissue Location

Expressed in brain > heart > skeletal muscle = kidney = pancreas = liver = placenta > lung. In brain, it is expressed in cerebellum, cerebral cortex, medulla, spinal cord, occipital lobe, frontal lobe, temporal lobe and putamen.

Images



Western blot analysis of ITPK1 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.