

Anti-ACAT1 Rabbit Monoclonal Antibody

Catalog # ABO16419

Product Information

Application	WB, IHC, IP
Primary Accession	P24752
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-ACAT1 Rabbit Monoclonal Antibody . Tested in WB, IHC, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	38
Other Names	Acetyl-CoA acetyltransferase, mitochondrial, 2.3.1.9 {ECO:0000255 PROSITE-ProRule:PRU10020, ECO:0000269 PubMed:1715688, ECO:0000269 PubMed:17371050, ECO:0000269 PubMed:7728148, ECO:0000269 PubMed:9744475}, Acetoacetyl-CoA thiolase, T2, ACAT1, ACAT, MAT
Calculated MW	45200
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 29A55
Immunogen	A synthesized peptide derived from human ACAT1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	ACAT1
-------------	-------

Synonyms

ACAT, MAT


Function

This is one of the enzymes that catalyzes the last step of the mitochondrial beta-oxidation pathway, an aerobic process breaking down fatty acids into acetyl-CoA (PubMed:[1715688](#), PubMed:[7728148](#), PubMed:[9744475](#)). Using free coenzyme A/CoA, catalyzes the thiolitic cleavage of medium- to long-chain 3-oxoacyl-CoAs into acetyl-CoA and a fatty acyl-CoA shortened by two carbon atoms (PubMed:[1715688](#), PubMed:[7728148](#), PubMed:[9744475](#)). The activity of the enzyme is reversible and it can also catalyze the condensation of two acetyl-CoA molecules into acetoacetyl-CoA (PubMed:[17371050](#)). Thereby, it plays a major role in ketone body metabolism (PubMed:[1715688](#), PubMed:[17371050](#), PubMed:[7728148](#), PubMed:[9744475](#)).


Cellular Location

Mitochondrion.

Images



Western blot analysis of ACAT1 expression in HepG2 cell lysate.



Immunoprecipitate (IP) analysis using the Antibody at 1:50 dilution. (wb at 1:1K dilution)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.