

Anti-MST4 Rabbit Monoclonal Antibody

Catalog # ABO16443

Product Information

Application	WB, IHC
Primary Accession	Q9P289
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-MST4 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	51765
Other Names	Serine/threonine-protein kinase 26, 2.7.11.1, MST3 and SOK1-related kinase, Mammalian STE20-like protein kinase 4, MST-4, STE20-like kinase MST4, Serine/threonine-protein kinase MASK, STK26 (HGNC:18174)
Calculated MW	46529
Application Details	WB 1:500-1:2000 IHC 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 29S79 A synthesized peptide derived from human MST4
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	STK26 (HGNC:18174)
Function	Serine/threonine-protein kinase that acts as a mediator of cell growth (PubMed: 11641781 , PubMed: 17360971). Modulates apoptosis (PubMed: 11641781 , PubMed: 17360971). In association with STK24 negatively

regulates Golgi reorientation in polarized cell migration upon RHO activation (PubMed:[27807006](#)). Phosphorylates ATG4B at 'Ser- 383', thereby increasing autophagic flux (PubMed:[29232556](#)). Part of the striatin-interacting phosphatase and kinase (STRIPAK) complexes. STRIPAK complexes have critical roles in protein (de)phosphorylation and are regulators of multiple signaling pathways including Hippo, MAPK, nuclear receptor and cytoskeleton remodeling. Different types of STRIPAK complexes are involved in a variety of biological processes such as cell growth, differentiation, apoptosis, metabolism and immune regulation (PubMed:[18782753](#)).

Cellular Location

Cytoplasm. Golgi apparatus Note=Colocalized with RIPOR1 in the Golgi of serum-starved cells and relocated to cytoplasmic punctae, probably vesicular compartments, along with RIPOR1 upon serum stimulation in a Rho- and PDCD10-dependent manner (PubMed:27807006).

Images



Western blot analysis of MST4 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.