

Anti-GNE Rabbit Monoclonal Antibody

Catalog # ABO16460

Product Information

Application	WB, IF, ICC
Primary Accession	Q9Y223
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-GNE Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human.

Additional Information

Gene ID	10020
Other Names	Bifunctional UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase, UDP-GlcNAc-2-epimerase/ManAc kinase, UDP-N-acetylglucosamine 2-epimerase (hydrolyzing), 3.2.1.183, UDP-GlcNAc-2-epimerase, Uridine diphosphate-N-acetylglucosamine-2-epimerase, N-acetylmannosamine kinase, 2.7.1.60, ManAc kinase, GNE (HGNC:23657)
Calculated MW	79275
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 29G96
Immunogen	A synthesized peptide derived from human GNE
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	GNE (HGNC:23657)
Function	Bifunctional enzyme that possesses both UDP-N- acetylglucosamine

2-epimerase and N-acetylmannosamine kinase activities, and serves as the initiator of the biosynthetic pathway leading to the production of N-acetylneuraminic acid (NeuAc), a critical precursor in the synthesis of sialic acids. By catalyzing this pivotal and rate-limiting step in sialic acid biosynthesis, this enzyme assumes a pivotal role in governing the regulation of cell surface sialylation, playing a role in embryonic angiogenesis (PubMed:[10334995](#), PubMed:[11326336](#), PubMed:[14707127](#), PubMed:[16503651](#), PubMed:[2808337](#), PubMed:[38237079](#)). Sialic acids represent a category of negatively charged sugars that reside on the surface of cells as terminal components of glycoconjugates and mediate important functions in various cellular processes, including cell adhesion, signal transduction, and cellular recognition (PubMed:[10334995](#), PubMed:[14707127](#)).


Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:O35826}

Tissue Location

Highest expression in liver and placenta. Also found in heart, brain, lung, kidney, skeletal muscle and pancreas Isoform 1 is expressed in heart, brain, kidney, liver, placenta, lung, spleen, pancreas, skeletal muscle and colon. Isoform 2 is expressed mainly in placenta, but also in brain, kidney, liver, lung, pancreas and colon. Isoform 3 is expressed at low level in kidney, liver, placenta and colon.

Images



Western blot analysis of GNE expression in K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.