

Anti-PCBP2 Rabbit Monoclonal Antibody

Catalog # ABO16484

Product Information

Application	WB, IF, ICC, IP
Primary Accession	Q15366
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-PCBP2 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	5094
Other Names	Poly(rC)-binding protein 2, Alpha-CP2, Heterogeneous nuclear ribonucleoprotein E2, hnRNP E2, PCBP2 {ECO:0000303 PubMed:7607214, ECO:0000312 HGNC:HGNC:8648}
Calculated MW	38580
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 30P20 A synthesized peptide derived from human PCBP2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	PCBP2 {ECO:0000303 PubMed:7607214, ECO:0000312 HGNC:HGNC:8648}
Function	Single-stranded nucleic acid binding protein that binds preferentially to oligo dC (PubMed: 12414943 , PubMed: 7607214). Major cellular poly(rC)-binding

protein (PubMed:[12414943](#)). Also binds poly(rU) (PubMed:[12414943](#)). Acts as a negative regulator of antiviral signaling (PubMed:[19881509](#), PubMed:[35322803](#)). Negatively regulates cellular antiviral responses mediated by MAVS signaling (PubMed:[19881509](#)). It acts as an adapter between MAVS and the E3 ubiquitin ligase ITCH, therefore triggering MAVS ubiquitination and degradation (PubMed:[19881509](#)). Negatively regulates the cGAS-STING pathway via interaction with CGAS, preventing the formation of liquid- like droplets in which CGAS is activated (PubMed:[35322803](#)). Together with PCBP1, required for erythropoiesis, possibly by regulating mRNA splicing (By similarity).


Cellular Location

Nucleus. Cytoplasm. Note=Loosely bound in the nucleus (PubMed:7607214). May shuttle between the nucleus and the cytoplasm (PubMed:7607214).


Tissue Location

Detected in all tissues examined.

Images



Western blot analysis of PCBP2 expression in HeLa cell lysate.



Immunoprecipitate (IP) analysis using the Antibody at 1:50 dilution. (wb at 1:1K dilution)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.