

Anti-SLC27A4 Rabbit Monoclonal Antibody

Catalog # ABO16535

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	Q6P1M0
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-SLC27A4 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human.

Additional Information

Gene ID	10999
Other Names	Long-chain fatty acid transport protein 4, FATP-4, Fatty acid transport protein 4, Arachidonate--CoA ligase, SLC27A4 (HGNC:10998)
Calculated MW	72064
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 30S71 A synthesized peptide derived from human SLC27A4
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	SLC27A4 (HGNC:10998)
Function	Mediates the levels of long-chain fatty acids (LCFA) in the cell by facilitating their transport across cell membranes (PubMed: 10518211 ,

PubMed:[12556534](#), PubMed:[20448275](#), PubMed:[21395585](#), PubMed:[22022213](#)). Appears to be the principal fatty acid transporter in small intestinal enterocytes (PubMed:[20448275](#)). Also functions as an acyl-CoA ligase catalyzing the ATP-dependent formation of fatty acyl- CoA using LCFA and very-long-chain fatty acids (VLCFA) as substrates, which prevents fatty acid efflux from cells and might drive more fatty acid uptake (PubMed:[22022213](#), PubMed:[24269233](#)). Plays a role in the formation of the epidermal barrier. Required for fat absorption in early embryogenesis (By similarity). Probably involved in fatty acid transport across the blood barrier (PubMed:[21395585](#)). Indirectly inhibits RPE65 via substrate competition and via production of VLCFA derivatives like lignoceroyl-CoA. Prevents light-induced degeneration of rods and cones (By similarity).


Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein


Tissue Location

Expressed at highest levels in brain, testis, colon and kidney. Expressed at medium levels in heart and liver, small intestine and stomach. Expressed at low levels in peripheral leukocytes, bone marrow, skeletal muscle and aorta. Expressed in adipose tissue (PubMed:[24269233](#), PubMed:[9878842](#)). Expressed in brain gray matter (PubMed:[21395585](#)).

Images



Western blot analysis of SLC27A4 expression in HepG2 cell lysate.



Immunoprecipitate (IP) analysis using the Antibody at 1:50 dilution. (wb at 1:1K dilution)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.