

Anti-GNAQ Rabbit Monoclonal Antibody

Catalog # ABO16538

Product Information

Application	WB
Primary Accession	P50148
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-GNAQ Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	2776
Other Names	Guanine nucleotide-binding protein G(q) subunit alpha, Guanine nucleotide-binding protein alpha-q, GNAQ, GAQ
Calculated MW	42142
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 30G74
Immunogen	A synthesized peptide derived from human GNAQ
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	GNAQ (HGNC:4390)
Synonyms	GAQ
Function	Guanine nucleotide-binding proteins (G proteins) function as transducers downstream of G protein-coupled receptors (GPCRs) in numerous signaling cascades (PubMed: 34556863 , PubMed: 35672283 , PubMed: 37991948). The

alpha chain contains the guanine nucleotide binding site and alternates between an active, GTP-bound state and an inactive, GDP-bound state (PubMed:[37991948](#)). Signaling by an activated GPCR promotes GDP release and GTP binding (PubMed:[37991948](#)). The alpha subunit has a low GTPase activity that converts bound GTP to GDP, thereby terminating the signal (PubMed:[37991948](#)). Both GDP release and GTP hydrolysis are modulated by numerous regulatory proteins (PubMed:[37991948](#)). Signaling is mediated via phospholipase C-beta- dependent inositol lipid hydrolysis for signal propagation: activates phospholipase C-beta: following GPCR activation, GNAQ activates PLC- beta (PLCB1, PLCB2, PLCB3 or PLCB4), leading to production of diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) (PubMed:[37991948](#)). Required for platelet activation (By similarity). Regulates B-cell selection and survival and is required to prevent B- cell-dependent autoimmunity (By similarity). Regulates chemotaxis of BM-derived neutrophils and dendritic cells (in vitro) (By similarity). Transduces FFAR4 signaling in response to long-chain fatty acids (LCFAs) (PubMed:[27852822](#)). Together with GNA11, required for heart development (By similarity).


Cellular Location

Cell membrane; Peripheral membrane protein; Cytoplasmic side. Golgi apparatus. Nucleus {ECO:0000250|UniProtKB:P21279} Nucleus membrane {ECO:0000250|UniProtKB:P21279}. Note=Associated to the plasma membrane via lipid-anchor (PubMed:19001095). Colocalizes with the adrenergic receptors ADRA1A and ADRA1B at the nuclear membrane of cardiac myocytes (By similarity). {ECO:0000250|UniProtKB:P21279, ECO:0000269|PubMed:19001095}


Tissue Location

Predominantly expressed in ovary, prostate, testis and colon. Down-regulated in the peripheral blood lymphocytes (PBLs) of rheumatoid arthritis patients (at protein level)


Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of GNAQ expression in (1) HeLa cell lysate; (2) NIH/3T3 cell lysate.



Immunofluorescent analysis using the Antibody at 1:50 dilution.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.