

Anti-PREPL Rabbit Monoclonal Antibody

Catalog # ABO16542

Product Information

Application	WB, IF, ICC, IP, FC
Primary Accession	Q4J6C6
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-PREPL Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	9581
Other Names	Prolyl endopeptidase-like, 3.4.21.-, Prolylendopeptidase-like, PREPL, KIAA0436
Calculated MW	83927
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 30P78
Immunogen	A synthesized peptide derived from human PREPL
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	PREPL
Synonyms	KIAA0436
Function	Serine peptidase whose precise substrate specificity remains unclear

(PubMed:[16143824](#), PubMed:[16385448](#), PubMed:[28726805](#)). Does not cleave peptides after a arginine or lysine residue (PubMed:[16143824](#)). Regulates trans-Golgi network morphology and sorting by regulating the membrane binding of the AP-1 complex (PubMed:[23321636](#)). May play a role in the regulation of synaptic vesicle exocytosis (PubMed:[24610330](#)).


Cellular Location

Cytoplasm, cytosol. Golgi apparatus, trans-Golgi network {ECO:0000250|UniProtKB:Q8C167}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q8C167}. Golgi apparatus {ECO:0000250|UniProtKB:Q8C167}. Nucleus Note=Co-localizes with AP-1 in the trans-Golgi network (By similarity) Co-localizes with MAP2 and ACTB on the cytoskeleton (By similarity) Co-localizes with STX6 and GOSR2 at the Golgi apparatus (By similarity). {ECO:0000250|UniProtKB:Q8C167}


Tissue Location

Expressed in pyramidal neurons of the temporal cortex and neocortex (at protein level) (PubMed:23485813). Widely expressed (PubMed:15913950, PubMed:16385448). Expressed at higher level in brain, skeletal muscle, heart and kidney (PubMed:15913950, PubMed:16385448). Expressed at the endplates in the neuromuscular junction (PubMed:24610330).

Images



Western blot analysis of PREPL expression in (1) MCF7 cell lysate; (2) Raw264.7 cell lysate.



Immunoprecipitate (IP) analysis using the Antibody at 1:50 dilution. (wb at 1:1K dilution)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.