

Anti-eIF3A Rabbit Monoclonal Antibody

Catalog # ABO16658

Product Information

Application	WB, IHC
Primary Accession	Q14152
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-eIF3A Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	8661
Other Names	Eukaryotic translation initiation factor 3 subunit A {ECO:0000255 HAMAP-Rule:MF_03000}, eIF3a {ECO:0000255 HAMAP-Rule:MF_03000}, Eukaryotic translation initiation factor 3 subunit 10 {ECO:0000255 HAMAP-Rule:MF_03000}, eIF-3-theta {ECO:0000255 HAMAP-Rule:MF_03000}, eIF3 p167, eIF3 p180, eIF3 p185, EIF3A {ECO:0000255 HAMAP-Rule:MF_03000}
Calculated MW	166569
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 31E29
Immunogen	A synthesized peptide derived from human eIF3A
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	EIF3A {ECO:0000255 HAMAP-Rule:MF_03000}
-------------	---


Function

RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:[17581632](#), PubMed:[25849773](#)). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:[11169732](#), PubMed:[17581632](#)). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem- loop binding to exert either translational activation or repression (PubMed:[25849773](#), PubMed:[27462815](#)).


Cellular Location

Cytoplasm {ECO:0000255 | HAMAP-Rule:MF_03000, ECO:0000269 | PubMed:9150439}


Images



Immunofluorescent analysis using the Antibody at 1:50 dilution.





Immunofluorescent analysis using the Antibody at 1:50 dilution.



Immunofluorescent analysis using the Antibody at 1:150 dilution.

Immunofluorescent analysis using the Antibody at 1:500 dilution.



Western blot analysis of eIF3A expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.