

Anti-HSPC150 Rabbit Monoclonal Antibody

Catalog # ABO16677

Product Information

Application	WB
Primary Accession	Q9NPD8
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-HSPC150 Rabbit Monoclonal Antibody . Tested in WB applications. This antibody reacts with Human.

Additional Information

Gene ID	29089
Other Names	Ubiquitin-conjugating enzyme E2 T, 2.3.2.23, Cell proliferation-inducing gene 50 protein, E2 ubiquitin-conjugating enzyme T, Ubiquitin carrier protein T, Ubiquitin-protein ligase T, UBE2T
Calculated MW	22521
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 31U48
Immunogen	A synthesized peptide derived from human HSPC150
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	UBE2T
Function	Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. Catalyzes monoubiquitination. Involved in mitomycin-C (MMC)-induced DNA repair. Acts as a specific E2 ubiquitin-conjugating enzyme for the Fanconi anemia complex by associating with E3 ubiquitin-protein

ligase FANCL and catalyzing monoubiquitination of FANCD2, a key step in the DNA damage pathway (PubMed:[16916645](#), PubMed:[17938197](#), PubMed:[19111657](#), PubMed:[19589784](#), PubMed:[28437106](#)). Also mediates monoubiquitination of FANCL and FANCI (PubMed:[16916645](#), PubMed:[17938197](#), PubMed:[19111657](#), PubMed:[19589784](#)). May contribute to ubiquitination and degradation of BRCA1 (PubMed:[19887602](#)). In vitro able to promote polyubiquitination using all 7 ubiquitin Lys residues, but may prefer 'Lys-11', 'Lys-27', 'Lys-48' and 'Lys-63'-linked polyubiquitination (PubMed:[20061386](#)).


Cellular Location

Nucleus. Note=Accumulates to chromatin

Images



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of HSPC150 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.