

# Anti-Myeloperoxidase Rabbit Monoclonal Antibody

Catalog # ABO16742

## Product Information

|                   |                                                                                                                                |
|-------------------|--------------------------------------------------------------------------------------------------------------------------------|
| Application       | WB, IHC                                                                                                                        |
| Primary Accession | <a href="#">P05164</a>                                                                                                         |
| Host              | Rabbit                                                                                                                         |
| Isotype           | Rabbit IgG                                                                                                                     |
| Reactivity        | Rat, Human, Mouse                                                                                                              |
| Clonality         | Monoclonal                                                                                                                     |
| Format            | Liquid                                                                                                                         |
| Description       | Anti-Myeloperoxidase Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat. |

## Additional Information

|                     |                                                                                                                                                                                              |
|---------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Gene ID             | 4353                                                                                                                                                                                         |
| Other Names         | Myeloperoxidase, MPO, 1.11.2.2, Myeloperoxidase, 89 kDa myeloperoxidase, 84 kDa myeloperoxidase, Myeloperoxidase light chain, Myeloperoxidase heavy chain, MPO ( <a href="#">HGNC:7218</a> ) |
| Calculated MW       | 83869                                                                                                                                                                                        |
| Application Details | WB 1:500-1:2000<br>IHC 1:50-1:200                                                                                                                                                            |
| Source              | Eukaryota                                                                                                                                                                                    |
| Contents            | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.                                                                          |
| Clone Names         | Clone: 32M13                                                                                                                                                                                 |
| Immunogen           | A synthesized peptide derived from human Myeloperoxidase                                                                                                                                     |
| Purification        | Affinity-chromatography                                                                                                                                                                      |
| Storage             | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.                                                   |

## Protein Information

|          |                                                                                                                                                                                                                                           |
|----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Name     | MPO {ECO:0000303 PubMed:21880720, ECO:0000312 HGNC:HGNC:7218}                                                                                                                                                                             |
| Function | Peroxidase that plays a central role in the host defense system of polymorphonuclear leukocytes by mediating both (1) formation of neutrophil extracellular trap (NETs) and (2) microbicidal activity (PubMed: <a href="#">16125131</a> , |

PubMed:[17438335](#), PubMed:[20974672](#), PubMed:[20974816](#), PubMed:[22131345](#), PubMed:[25066128](#), PubMed:[28574339](#), PubMed:[40963017](#), PubMed:[4978299](#), PubMed:[6295491](#)). Promotes NET formation by mediating chromatin disassembly: translocates to the nucleus and specifically binds to nucleosomes, both as monomer and homodimer, leading to nucleosome unstacking and initial chromatin decondensation (PubMed:[40963017](#)). Homodimers clash with one end of the nucleosomal DNA, leading to DNA unwrapping, initiating complete disassembly of nucleosomes and chromatin transformation into NETs in an ATP-independent manner (PubMed:[40963017](#)). NETs, which are mainly composed of DNA fibers and globular proteins, are then extruded into the extracellular space by neutrophils to trap pathogens and release antimicrobial proteins to destroy them (PubMed:[40963017](#)). Participates to the microbicidal activity against a wide range of organisms by acting as a peroxidase that catalyzes the formation of oxidants in presence of hydrogen peroxide (PubMed:[16125131](#), PubMed:[17074761](#), PubMed:[17438335](#), PubMed:[21880720](#), PubMed:[22131345](#), PubMed:[22352991](#), PubMed:[22902565](#), PubMed:[24194519](#), PubMed:[33273015](#), PubMed:[9359420](#), PubMed:[9922160](#)). Mediates the formation of hypohalous acids, mainly hypochlorous acid (HOCl) in physiologic situations, that greatly enhance polymorphonuclear leukocyte microbicidal activity (PubMed:[16125131](#), PubMed:[17074761](#), PubMed:[17438335](#), PubMed:[19608745](#), PubMed:[21880720](#), PubMed:[22902565](#), PubMed:[24194519](#), PubMed:[9359420](#), PubMed:[9922160](#)). In addition to hypochlorous acid, catalyzes formation of hypobromous acid (HOBr), hypoiodous acid (HOI) and hypothiocyanous acid (HOSCN) (PubMed:[1318692](#), PubMed:[16125131](#), PubMed:[17438335](#), PubMed:[19608745](#), PubMed:[6295491](#), PubMed:[9359420](#)). Also catalyzes oxidation of nitrite into the highly reactive nitrogen dioxide radical (PubMed:[10777476](#), PubMed:[8385644](#), PubMed:[9450756](#)). Formation of oxidants are widely believed to be responsible for much of the anti-bactericidal activity of neutrophils (PubMed:[8385644](#), PubMed:[9450756](#)). Oxidants, such as hypochlorous acid or nitrogen dioxide radical, can also oxidize amino acid residues on proteins and generate chlorination and nitration post-translational modifications, respectively (PubMed:[15314690](#), PubMed:[15326314](#), PubMed:[15498770](#), PubMed:[15574409](#), PubMed:[16091367](#), PubMed:[7622459](#)). Chlorination and nitration of the lipid-free form of APOA1 impairs cholesterol transport (PubMed:[15326314](#), PubMed:[15498770](#), PubMed:[15574409](#)). Superoxides generated by MPO can also promote dioxygenation of tryptophan residues on proteins (PubMed:[16091367](#), PubMed:[40081572](#)). Also able to oxidize melatonin into N1-acetyl-N2-formyl-5-methoxykynuramine either in presence of hydrogen peroxide or superoxide (PubMed:[15636586](#), PubMed:[16148002](#)). Oxidizes urate into 5-hydroxyisourate (PubMed:[21266577](#)). Functions as a nitric oxide (NO) oxidase during inflammation, by catalytically consuming NO, impairing NO's ability to maintain vascular tone and function (PubMed:[12089442](#)). May also mediate the proteolytic cleavage of alpha-1-microglobulin to form t-alpha-1- microglobulin, which potently inhibits oxidation of low-density lipoprotein particles and limits vascular damage (PubMed:[25698971](#)).

## Cellular Location


Lysosome. Cytoytic granule. Cytoplasmic vesicle, phagosome. Secreted. Nucleus. Chromosome Note=Localizes to azurophil granules, a type of primary lysosome found in neutrophils and monocytes that contains antimicrobial proteins (PubMed:[25066128](#), PubMed:[2981589](#), PubMed:[6095812](#), PubMed:[6192815](#)). A substantial fraction is also constitutively secreted by normal bone marrow granulocyte precursors (PubMed:[6095812](#)). During formation of neutrophil extracellular traps (NETs), translocates into the nucleus and associates to chromatin, leading to chromatin dislocation and its transformation into NETs (PubMed:[25066128](#), PubMed:[40963017](#)). Core component of NETs (PubMed:[22131345](#), PubMed:[32434865](#), PubMed:[40963017](#)).

## Tissue Location

Specifically expressed in neutrophils and monocytes.

## Images

---



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.