

Anti-VAPA Rabbit Monoclonal Antibody

Catalog # ABO16758

Product Information

Application	WB, IHC
Primary Accession	Q9P0L0
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-VAPA Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	9218
Other Names	Vesicle-associated membrane protein-associated protein A, VAMP-A, VAMP-associated protein A, VAP-A, 33 kDa VAMP-associated protein {ECO:0000303 Ref.2}, VAP-33, VAPA (HGNC:12648), VAP33
Calculated MW	27893
Application Details	WB 1:500-1:2000 IHC 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 32V29 A synthesized peptide derived from human VAPA
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	VAPA (HGNC:12648)
Synonyms	VAP33
Function	Endoplasmic reticulum (ER)-anchored protein that mediates the formation of

contact sites between the ER and endosomes via interaction with FFAT motif-containing proteins such as STARD3 or WDR44 (PubMed:[32344433](#), PubMed:[33124732](#)). STARD3-VAPA interaction enables cholesterol transfer from the ER to endosomes (PubMed:[33124732](#)). Via interaction with WDR44 participates in neosynthesized protein export (PubMed:[32344433](#)). In addition, recruited to the plasma membrane through OSBPL3 binding (PubMed:[25447204](#)). The OSBPL3-VAPA complex stimulates RRAS signaling which in turn attenuates integrin beta-1 (ITGB1) activation at the cell surface (PubMed:[25447204](#)). With OSBPL3, may regulate ER morphology (PubMed:[16143324](#)). May play a role in vesicle trafficking (PubMed:[11511104](#), PubMed:[19289470](#)).


Cellular Location

Endoplasmic reticulum membrane; Single-pass type IV membrane protein. Cell membrane; Single-pass type IV membrane protein. Cell junction, tight junction. Nucleus membrane {ECO:0000250|UniProtKB:Q9Z270}. Note=Present in the plasma membrane and in intracellular vesicles, together with SNARE proteins. May also associate with the cytoskeleton. Colocalizes with OCLN at the tight junction in polarized epithelial cells.


Tissue Location

Ubiquitous.

Images



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.