

Anti-SYVN1 / HRD1 Rabbit Monoclonal Antibody

Catalog # ABO16767

Product Information

Application	WB, IP, FC
Primary Accession	Q86TM6
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-SYVN1 / HRD1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	84447
Other Names	E3 ubiquitin-protein ligase synoviolin, 2.3.2.27, RING-type E3 ubiquitin transferase synoviolin, Synovial apoptosis inhibitor 1, SYVN1, HRD1, KIAA1810
Calculated MW	67685
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 32S38 A synthesized peptide derived from human SYVN1 / HRD1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	SYVN1 {ECO:0000303 PubMed:15489334}
Function	E3 ubiquitin-protein ligase which accepts ubiquitin specifically from endoplasmic reticulum-associated UBC7 E2 ligase and transfers it to

substrates, promoting their degradation (PubMed:[12459480](#), PubMed:[12646171](#), PubMed:[12975321](#), PubMed:[14593114](#), PubMed:[16289116](#), PubMed:[16847254](#), PubMed:[17059562](#), PubMed:[17141218](#), PubMed:[17170702](#), PubMed:[22607976](#), PubMed:[27827840](#), PubMed:[26471130](#), PubMed:[28827405](#)). Component of the endoplasmic reticulum quality control (ERQC) system also called ER-associated degradation (ERAD) involved in ubiquitin-dependent degradation of misfolded endoplasmic reticulum proteins (PubMed:[12459480](#), PubMed:[12646171](#), PubMed:[12975321](#), PubMed:[14593114](#), PubMed:[16289116](#), PubMed:[16847254](#), PubMed:[17059562](#), PubMed:[17141218](#), PubMed:[17170702](#), PubMed:[22607976](#), PubMed:[26471130](#), PubMed:[28842558](#)). Also promotes the degradation of normal but naturally short-lived proteins such as SGK. Protects cells from ER stress-induced apoptosis. Protects neurons from apoptosis induced by polyglutamine- expanded huntingtin (HTT) or unfolded GPR37 by promoting their degradation (PubMed:[17141218](#)). Sequesters p53/TP53 in the cytoplasm and promotes its degradation, thereby negatively regulating its biological function in transcription, cell cycle regulation and apoptosis (PubMed:[17170702](#)). Mediates the ubiquitination and subsequent degradation of cytoplasmic NFE2L1 (By similarity). During the early stage of B cell development, required for degradation of the pre-B cell receptor (pre-BCR) complex, hence supporting further differentiation into mature B cells (By similarity).


Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein


Tissue Location

Ubiquitously expressed, with highest levels in liver and kidney (at protein level). Up-regulated in synovial tissues from patients with rheumatoid arthritis (at protein level)

Images



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Immunoprecipitate (IP) analysis using the Antibody at 1:50 dilution. (wb at 1:1K dilution)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.