

GITR / TNFSF18 Polyclonal Antibody

Rabbit Polyclonal Antibody Catalog # ABV11779

Product Information

| Application WB | |
|--------------------------------|-----------|
| Primary Accession <u>Q9UNC</u> | <u>52</u> |
| Reactivity Humar | 1 |
| Host Rabbit | |
| Clonality Polyclo | nal |
| Isotype Rabbit | IgG |
| Calculated MW 20308 | |

Additional Information

| Gene ID | 8995 |
|--|---|
| Positive Control Application & Usage Alias Symbol Other Names | WB WB: 1:1000 TNFRSF18 Tumor necrosis factor ligand superfamily member 18, Activation-inducible TNF-related ligand, AITRL, Glucocorticoid-induced TNF-related ligand, hGITRL, TNFSF18, AITRL, GITRL, TL6 |
| Appearance | Colorless liquid |
| Formulation | PBS with 0.09% (W/V) sodium azide. |
| Reconstitution & Storage | -20 °C |
| Background Descriptions Precautions | GITR / TNFSF18 Polyclonal Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| Name | TNFSF18 (<u>HGNC:11932</u>) |
|----------|--|
| Synonyms | AITRL, GITRL, TL6 |
| Function | Cytokine that binds to TNFRSF18/AITR/GITR. Regulates T-cell responses. Can function as costimulator and lower the threshold for T- cell activation and T-cell proliferation. Important for interactions between activated T-lymphocytes and endothelial cells. Mediates activation of NF-kappa-B. Triggers increased phosphorylation of STAT1 and up-regulates expression of VCAM1 and ICAM1 (PubMed: <u>23892569</u>). Promotes leukocyte adhesion to |

| | endothelial cells (PubMed: <u>23892569</u>). Regulates migration of monocytes from the splenic reservoir to sites of inflammation (By similarity). |
|-------------------|---|
| Cellular Location | Cell membrane; Single-pass type II membrane protein |
| Tissue Location | Expressed at high levels in the small intestine, ovary, testis, kidney and endothelial cells |

Background

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for receptor TNFRSF18/AITR/GITR. It has been shown to modulate T lymphocyte survival in peripheral tissues. This cytokine is also found to be expressed in endothelial cells, and is thought to be important for interaction between T lymphocytes and endothelial cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.