

Lactoferrin Antibody

Rabbit Polyclonal Antibody

Catalog # ABV11802

Product Information

Application	WB
Primary Accession	P02788
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	78182

Additional Information

Gene ID	4057
Positive Control	WB: human lactoferrin
Application & Usage	WB: 1-4 μ g
Alias Symbol	LTF
Other Names	Growth-inhibiting protein 12, Talalactoferrin, Lactotransferrin
Appearance	Colorless liquid
Formulation	In PBS pH 7.2, 0.01 % BSA, 0.03 % ProClin® and 50 % glycerol
Reconstitution & Storage	-20 °C
Background Descriptions	
Precautions	Lactoferrin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	LTF (HGNC:6720)
Synonyms	GIG12, LF
Function	Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate.
Cellular Location	[Isoform 1]: Secreted. Cytoplasmic granule. Note=Secreted into most exocrine fluids by various endothelial cells Stored in the secondary granules of neutrophils
Tissue Location	High levels are found in saliva and tears, intermediate levels in serum and

plasma, and low levels in urine. In kidney, detected in the distal collecting tubules in the medulla but not in the cortical region or in blood vessels. Detected in peripheral blood neutrophils (at protein level). Isoform 1 and isoform DeltaLf are expressed in breast, prostate, spleen, pancreas, kidney, small intestine, lung, skeletal muscle, uterus, thymus and fetal liver Isoform 1 is expressed in brain, testis and peripheral blood leukocytes; isoform DeltaLf is barely detectable in these tissues Isoform DeltaLf is expressed in placenta, liver and ovary; isoform 1 is barely detectable in these tissues. In kidney, isoform 1 is expressed at high levels in the collecting tubules of the medulla but at very low levels in the cortex.

Background

Lactoferrin is a glycoprotein present in exocrine secretions and in the secondary granules of polymorphonuclear neutrophils. It is the iron binding protein in milk. Lactoferrin is known to regulate intestinal iron absorption, and to participate in the defense against bacteria.

Images



Western blot with human Lactoferrin antibody in: Lane1: h Lactoferrin, 2ng; Lane2: h Lactoferrin, 10ng; Lane3: h Lactoferrin, 50ng

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.