10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Phospho-BID (Ser78) Antibody

Rabbit Polyclonal Antibody
Catalog # ABV11846

Product Information

Application	WB, IHC
Primary Accession	P55957
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	21995

Additional Information

Gene ID	637
Positive Control	WB: mouse spleen, rat spleen lysates; IHC: human brain tissue
Application & Usage	WB; 1:500 – 1:2000, IHC; 1:50 – 1:200
Alias Symbol	BID
Other Names	BH3-interacting domain death agonist, p22 BID, BID
Appearance	Colorless liquid
Formulation	In 0.42% Potassium phosphate; 0.87% Sodium chloride; pH 7.3; 30% glycerol and 0.01% sodium azide
Reconstitution & Storage	-20 °C
Background Descriptions	
Precautions	Phospho-BID (Ser78) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	BID
Function	Induces caspases and apoptosis (PubMed: 14583606). Counters the protective effect of BCL2 (By similarity).
Cellular Location	Cytoplasm. Mitochondrion membrane. Mitochondrion outer membrane. Note=When uncleaved, it is predominantly cytoplasmic. [BH3-interacting domain death agonist p13]: Mitochondrion membrane {ECO:0000250|UniProtKB:P70444}. Note=Associated with the mitochondrial membrane. {ECO:0000250|UniProtKB:P70444} [Isoform 3]: Cytoplasm

Tissue Location	[Isoform 2]: Expressed in spleen, pancreas and placenta (at protein level). [Isoform 4]: Expressed in lung and pancreas (at protein level).


Background

Bid, a BH3 domain-containing proapoptotic Bcl-2 family member, is localized in the cytosolic fraction of cells as an inactive precursor. Its active form is generated upon proteolytic cleavage by caspase-8 in the Fas signaling pathway. Cleaved Bid translocates to mitochondria and releases its potent proapoptotic activity, which in turn induces cytochrome c release and mitochondrial damage. The cytochrome c releasing activity of Bid was antagonized by Bcl-2. Mutation in the SH3 domain can diminish the cytochrome c releasing activity. In the animal model studies, Bid-deficient mice are found resistant to the lethal effects of death factor signals relayed through Fas.

Images



Immunohistochemical analysis of BID(pS78) staining in H.brain formalin fixed paraffin embedded tissue section.



Western blot analysis of BID(pS78) expression in M.spleen(A), R.spleen(B) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.