

Anti-MARCH2 Antibody

Rabbit Polyclonal antibody

Catalog # ABV11872

Product Information

Application	WB, IHC, IF
Primary Accession	Q9P0N8
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	26995

Additional Information

Gene ID	51257
Positive Control	WB: HeLa, PC12, cell lysate; IHC: human brain tissue; IFC: HeLa cells
Application & Usage	WB; 1:500 – 1:2000, IHC; 1:50 – 1:200, IF/IC; 1:50 – 1:100
Alias Symbol	MARCH2
Other Names	RNF172, E3 ubiquitin-protein ligase MARCH2, Membrane-associated RING finger protein 2, Membrane-associated RING-CH protein II, MARCH-II, RING finger protein 172
Formulation	In 0.42% Potassium phosphate; 0.87% Sodium chloride; pH 7.3; 30% glycerol and 0.01% sodium azide
Reconstitution & Storage	12 months under -20°C
Background Descriptions	
Precautions	Anti-MARCH2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MARCHF2 (HGNC:28038)
Synonyms	MARCH2, RNF172
Function	E3 ubiquitin-protein ligase that may mediate ubiquitination of TFRC and CD86, and promote their subsequent endocytosis and sorting to lysosomes via multivesicular bodies. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates (PubMed: 14722266 , PubMed: 16428329). Together with GOPC/CAL mediates the ubiquitination and lysosomal degradation of CFTR (PubMed: 23818989). Ubiquitinates and

therefore mediates the degradation of DLG1 (PubMed:[17980554](#)). Regulates the intracellular trafficking and secretion of alpha1-antitrypsin/SERPINA1 and HP/haptoglobin via ubiquitination and degradation of the cargo receptor ERGIC3 (PubMed:[31142615](#)). Negatively regulates the antiviral and antibacterial immune response by repression of the NF-kB and type 1 IFN signaling pathways, via MARCHF2-mediated K48-linked polyubiquitination of IKBKG/NEMO, resulting in its proteasomal degradation (PubMed:[32935379](#)). May be involved in endosomal trafficking through interaction with STX6 (PubMed:[15689499](#)).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q5I0I2}. Lysosome membrane; Multi-pass membrane protein. Endosome membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q5I0I2}. Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasm. Cell membrane; Multi-pass membrane protein


Tissue Location

Broadly expressed..


Background

E3 ubiquitin-protein ligase that may mediate ubiquitination of TFRC and CD86, and promote their subsequent endocytosis and sorting to lysosomes via multivesicular bodies. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates. May be involved in endosomal trafficking through interaction with STX6

Images




Immunohistochemical analysis of MARCH2 staining in human brain formalin fixed paraffin embedded tissue section.



Immunofluorescent analysis of MARCH2 staining in HeLa cells.

Western blot analysis of MARCH2 expression in HeLa(A), PC12(B) whole cell lysates.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.