

Anti-EEF1A Antibody

Catalog # ABV11938

Product Information

Application	WB
Primary Accession	P68104
Reactivity	Human, Mouse, Rat
Host	Rabbit
Isotype	Rabbit IgG
Calculated MW	50141

Additional Information

Gene ID	1915
Positive Control	WB: H9C2 PMA-treated (A) whole cell lysates
Application & Usage	WB; 1:500 – 1:2000
Other Names	AcK146, EEF1A1, EEF1A, EF1A, LENG7, Elongation factor 1-alpha 1, EF-1-alpha-1, Elongation factor Tu, EF-Tu; Eukaryotic elongation factor 1 A-1, eEF1A-1, Leukocyte receptor cluster member 7, EEF1A2, EEF1AL, STN, Elongation factor 1-alpha 2, EF-1-alpha-2, Eukaryotic elongation factor 1 A-2, eEF1A-2, Statin-S1, EEF1A1P5, EEF1AL3, Putative elongation factor 1-alpha-like 3, EF-1-alpha-like 3, Eukaryotic elongation factor 1 A-like 3, eEF1A-like 3, Eukaryotic translation elongation factor 1 alpha-1 pseudogene 5
Target/Specificity	EEF1A1
Antibody Form	Liquid
Appearance	Colorless liquid
Handling	The antibody solution should be gently mixed before use
Reconstitution & Storage	-20°C
Background Descriptions	
Precautions	Anti-EEF1A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	EEF1A1
Synonyms	EEF1A, EF1A, LENG7
Function	Translation elongation factor that catalyzes the GTP- dependent binding of aminoacyl-tRNA (aa-tRNA) to the A-site of ribosomes during the elongation

phase of protein synthesis (PubMed:[26593721](#), PubMed:[26651998](#), PubMed:[36123449](#), PubMed:[36264623](#), PubMed:[36638793](#)). Base pairing between the mRNA codon and the aa-tRNA anticodon promotes GTP hydrolysis, releasing the aa-tRNA from EEF1A1 and allowing its accommodation into the ribosome (PubMed:[26593721](#), PubMed:[26651998](#), PubMed:[36123449](#), PubMed:[36264623](#), PubMed:[36638793](#)). The growing protein chain is subsequently transferred from the P-site peptidyl tRNA to the A-site aa-tRNA, extending it by one amino acid through ribosome-catalyzed peptide bond formation (PubMed:[26593721](#), PubMed:[26651998](#), PubMed:[36123449](#), PubMed:[36264623](#)). Also plays a role in the positive regulation of IFNG transcription in T-helper 1 cells as part of an IFNG promoter-binding complex with TXK and PARP1 (PubMed:[17177976](#)). Also plays a role in cytoskeleton organization by promoting actin bundling (By similarity).

Cellular Location

Cytoplasm. Nucleus. Nucleus, nucleolus. Cell membrane. Note=Colocalizes with DLC1 at actin-rich regions in the cell periphery (PubMed:19158340). Translocates together with ZPR1 from the cytoplasm to the nucleus and nucleolus after treatment with mitogens (PubMed:8650580). Localization at the cell membrane depends on EEF1A1 phosphorylation status and the presence of PPP1R16B (PubMed:26497934).

Images



WB analysis of EEF1A (Ack146) expression in H9C2 PMA_ treated (A) whole cell lysates

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.