

# Anti-MEF2A Antibody

Catalog # ABV11942

## Product Information

---

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | WB                     |
| <b>Primary Accession</b> | <a href="#">Q02078</a> |
| <b>Reactivity</b>        | Human, Mouse, Rat      |
| <b>Host</b>              | Rabbit                 |
| <b>Isotype</b>           | Rabbit IgG             |
| <b>Calculated MW</b>     | 54811                  |

## Additional Information

---

|                                     |   |
|-------------------------------------|---|
| <b>Gene ID</b>                      | 4205  |
| <b>Positive Control</b>             | WB; HEK293T, mouse muscle, rat muscle cell lysates  |
| <b>Application &amp; Usage</b>      | WB; 1:500 – 1:2000  |
| <b>Other Names</b>                  | MEF2; Myocyte-specific enhancer factor 2A; Serum response factor-like protein 1                       |
| <b>Target/Specificity</b>           | MEF2A   |
| <b>Antibody Form</b>                | Liquid  |
| <b>Appearance</b>                   | Colorless liquid  |
| <b>Handling</b>                     | The antibody solution should be gently mixed before use   |
| <b>Reconstitution &amp; Storage</b> | -20°C   |
| <b>Background Descriptions</b>      |   |
| <b>Precautions</b>                  | Anti-MEF2A Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

---

|                 |   |
|-----------------|---|
| <b>Name</b>     | MEF2A   |
| <b>Synonyms</b> | MEF2  |
| <b>Function</b> | Transcriptional activator which binds specifically to the MEF2 element, 5'-YTA[AT](4)TAR-3', found in numerous muscle-specific genes. Also involved in the activation of numerous growth factor- and stress-induced genes. Mediates cellular functions not only in skeletal and cardiac muscle development, but also in neuronal differentiation and survival. Plays diverse roles in the control of cell growth, survival and apoptosis via p38 MAPK signaling in muscle-specific and/or growth factor-related transcription. In |

cerebellar granule neurons, phosphorylated and sumoylated MEF2A represses transcription of NUR77 promoting synaptic differentiation. Associates with chromatin to the ZNF16 promoter.

**Cellular Location**


Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00251, ECO:0000269 | PubMed:12691662, ECO:0000269 | PubMed:16563226}

**Tissue Location**

Isoform MEF2 and isoform MEFA are expressed only in skeletal and cardiac muscle and in the brain. Isoform RSRFC4 and isoform RSRFC9 are expressed in all tissues examined

**Images**

---



WB analysis of MEF2A (AcK403) expression in HEK293T (A): mouse muscle (B): rat muscle (C) whole cell lysates

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.