

Anti-Pyruvate Dehydrogenase E2 Antibody (4A4-B6-C10)

Mouse Monoclonal Antibody

Catalog # ABV12053

Product Information

Application	WB, IF, IP
Primary Accession	P10515
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Clone Names	4A4-B6-C10
Calculated MW	68997

Additional Information

Gene ID	1737
Application & Usage Other Names	WB: Jurkat, A549, U251, F9, Lncap and Hela cell lysates, 70 kDa mitochondrial autoantigen of primary biliary cirrhosis, PBC, Dihydrolipoamide acetyltransferase component of pyruvate dehydrogenase complex, M2 antigen complex 70 kDa subunit, Pyruvate dehydrogenase complex component E2
Target/Specificity	Pyruvate Dehydrogenase E2
Antibody Form	Liquid
Appearance	Colorless liquid
Formulation	In buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.2% sodium azide, 50% glycerol
Handling	The antibody solution should be gently mixed before use.
Reconstitution & Storage	-20 °C
Background Descriptions Precautions	Anti-Pyruvate Dehydrogenase E2 Antibody (4A4-B6-C10) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DLAT (HGNC:2896)
Synonyms	DLTA

Function	As part of the pyruvate dehydrogenase complex, catalyzes the transfers of an acetyl group to a lipoic acid moiety (Probable). The pyruvate dehydrogenase complex, catalyzes the overall conversion of pyruvate to acetyl-CoA and CO(2), and thereby links cytoplasmic glycolysis and the mitochondrial tricarboxylic acid (TCA) cycle (Probable).
Cellular Location	Mitochondrion matrix {ECO:0000250 UniProtKB:P08461}

Background

The pyruvate dehydrogenase complex catalyzes the overall conversion of pyruvate to acetyl-CoA and CO₂, and thereby links the glycolytic pathway to the tricarboxylic cycle

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.