

Anti-GRP78/BiP Antibody (9E4-2A7-H6)

Mouse Monoclonal Antibody

Catalog # ABV12064

Product Information

Application	WB, IHC, IF
Primary Accession	P11021
Reactivity	Human, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Clone Names	9E4-2A7-H6
Calculated MW	72333

Additional Information

Gene ID	3309
Application & Usage	WB: HeLa, C6, Lncap and MDA-MB-468 cell lysates; IF: HeLa cells; IHC: Colorectal cancer tissue
Other Names	GRP-78, Endoplasmic reticulum luminal Ca(2+)-binding protein grp78, Heat shock 70 kDa protein 5, Immunoglobulin heavy chain-binding protein, BiP, HSPA5, GRP78
Target/Specificity	GRP78
Antibody Form	Liquid
Appearance	Colorless liquid
Formulation	In PBS (pH 7.4) containing with 0.02% sodium azide and 50% glycerol
Handling	The antibody solution should be gently mixed before use.
Reconstitution & Storage	-20 °C
Background Descriptions	
Precautions	Anti-GRP78/BiP Antibody (9E4-2A7-H6) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HSPA5 (HGNC:5238)
Function	Endoplasmic reticulum chaperone that plays a key role in protein folding and quality control in the endoplasmic reticulum lumen (PubMed: 2294010 ,

PubMed:[23769672](#), PubMed:[23990668](#), PubMed:[28332555](#)). Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10/ERdj5, probably to facilitate the release of DNAJC10/ERdj5 from its substrate (By similarity). Acts as a key repressor of the EIF2AK3/PERK and ERN1/IRE1- mediated unfolded protein response (UPR) (PubMed:[11907036](#), PubMed:[1550958](#), PubMed:[19538957](#), PubMed:[36739529](#)). In the unstressed endoplasmic reticulum, recruited by DNAJB9/ERdj4 to the luminal region of ERN1/IRE1, leading to disrupt the dimerization of ERN1/IRE1, thereby inactivating ERN1/IRE1 (By similarity). Also binds and inactivates EIF2AK3/PERK in unstressed cells (PubMed:[11907036](#)). Accumulation of misfolded protein in the endoplasmic reticulum causes release of HSPA5/BiP from ERN1/IRE1 and EIF2AK3/PERK, allowing their homodimerization and subsequent activation (PubMed:[11907036](#)). Plays an auxiliary role in post-translational transport of small presecretory proteins across endoplasmic reticulum (ER). May function as an allosteric modulator for SEC61 channel-forming translocon complex, likely cooperating with SEC62 to enable the productive insertion of these precursors into SEC61 channel. Appears to specifically regulate translocation of precursors having inhibitory residues in their mature region that weaken channel gating. May also play a role in apoptosis and cell proliferation (PubMed:[26045166](#)).


Cellular Location

Endoplasmic reticulum lumen. Melanosome. Cytoplasm {ECO:0000250|UniProtKB:P20029}. Cell surface Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545). Localizes to the cell surface of epithelial cells in response to high levels of free iron (PubMed:20484814, PubMed:24355926, PubMed:27159390)

Background



Probably plays a role in facilitating the assembly of multimeric protein complexes inside the endoplasmic reticulum. Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10, probably to facilitate the release of DNAJC10 from its substrate.

Images



Western blot detection of BiP/GRP78 (Cterminus) in HeLa, C6, Lncap and MOA-MB- 468 cell lysates using BiP/GRP78 (Cterminus) mouse mAb

Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using anti-BiP/GRP78 (C-terminus) mouse mAb



Immunohistochemical analysis of paraffinembedded
Colorectal cancer using BiP/GRP78 (C-terminus) Mouse
mAb

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.