10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CD19

Rabbit Monoclonal antibody(Mab)
Catalog # AD80005

Product Information

Application	IHC-P
Primary Accession	P15391
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Clone Names	391B5F3
Calculated MW	61128

Additional Information


Gene ID	930
Gene Name	CD19
Other Names	B-lymphocyte antigen CD19, B-lymphocyte surface antigen B4, Differentiation antigen CD19, T-cell surface antigen Leu-12, CD19, CD19
Dilution	IHC-P~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.
Precautions	CD19 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	CD19
Function	Functions as a coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes (PubMed:29523808). Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed:1373518, PubMed:16672701, PubMed:2463100). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed:12387743, PubMed:16672701, PubMed:9317126, PubMed:9382888). Is not required for early steps during B cell differentiation in the blood marrow (PubMed:9317126). Required for normal differentiation of B-1 cells (By similarity). Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed:1373518, PubMed:2463100). Required for normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed:12387743, PubMed:16672701, PubMed:9317126).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Membrane raft {ECO:0000250|UniProtKB:P25918}; Single-pass type I membrane protein

Tissue Location	{ECO:0000250|UniProtKB:P25918} Detected on marginal zone and germinal center B cells in lymph nodes (PubMed:2463100). Detected on blood B cells (at protein level) (PubMed:16672701, PubMed:2463100)


Images




Immunohistochemical analysis of formalin-fixed and paraffin-embedded human tonsil tissue was performed with CD19 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human tonsil tissue was performed with CD19 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human tonsil tissue was performed with CD19 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human tonsil tissue was performed with CD19 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.