



Desmin

Mouse Monoclonal antibody(Mab)
Catalog # AD80025

Product Information

Application IHC-P
Primary Accession P17661
Reactivity Human
Host Mouse
Clonality Monoclonal
Clone Names 105A3E2
Calculated MW 53536

Additional Information

Gene ID 1674

Other Names Desmin, DES

Dilution IHC-P~~Ready-to-use

Storage Maintain refrigerated at 2-8°C.

Protein Information

Name DES

Function Muscle-specific type III intermediate filament essential for proper muscular

structure and function. Plays a crucial role in maintaining the structure of sarcomeres, inter-connecting the Z-disks and forming the myofibrils, linking them not only to the sarcolemmal cytoskeleton, but also to the nucleus and mitochondria, thus providing strength for the muscle fiber during activity (PubMed: 25358400). In adult striated muscle they form a fibrous network connecting myofibrils to each other and to the plasma membrane from the periphery of the Z- line structures (PubMed:24200904, PubMed:25394388, PubMed: <u>26724190</u>). May act as a sarcomeric microtubule-anchoring protein: specifically associates with detyrosinated tubulin-alpha chains, leading to buckled microtubules and mechanical resistance to contraction. Required for nuclear membrane integrity, via anchoring at the cell tip and nuclear envelope, resulting in maintenance of microtubule-derived intracellular mechanical forces (By similarity). Contributes to the transcriptional regulation of the NKX2-5 gene in cardiac progenitor cells during a short period of cardiomyogenesis and in cardiac side population stem cells in the adult. Plays a role in maintaining an optimal conformation of nebulette (NEB) on heart muscle sarcomeres to bind and recruit cardiac alpha-actin (By similarity). Cytoplasm, myofibril, sarcomere, Z line. Cytoplasm Cell membrane,


sarcolemma. Nucleus {ECO:0000250 | UniProtKB:P31001}. Cell tip

Cellular Location

1 of 2

{ECO:0000250 | UniProtKB:P31001}. Nucleus envelope {ECO:0000250 | UniProtKB:P31001}. Note=Localizes in the intercalated disks which occur at the Z line of cardiomyocytes (PubMed:24200904, PubMed:26724190). Localizes in the nucleus exclusively in differentiating cardiac progenitor cells and premature cardiomyocytes (By similarity). PKP2 is required for correct anchoring of DES at the cell tip and nuclear envelope (By similarity) {ECO:0000250 | UniProtKB:P31001, ECO:0000269 | PubMed:24200904, ECO:0000269 | PubMed:26724190}

Images



鼠单抗

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.