10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# PCNA

Rabbit Monoclonal antibody(Mab)
Catalog # AD80035

## Product Information

| | |
|---|---|
| **Application** | IHC-P |
| **Primary Accession** | P12004 |
| **Reactivity** | Human |
| **Host** | Rabbit |
| **Clonality** | Monoclonal |
| **Clone Names** | 111B2E9 |
| **Calculated MW** | 28769 |

## Additional Information


| | |
|---|---|
| **Gene ID** | 5111 |
| **Gene Name** | PCNA |
| **Other Names** | Proliferating cell nuclear antigen, PCNA, Cyclin, PCNA |
| **Dilution** | IHC-P~~Ready-to-use |
| **Storage** | Maintain refrigerated at 2-8°C. |
| **Precautions** | PCNA Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | PCNA |
| **Function** | Confers DNA tethering and processivity to DNA polymerases and other proteins (PubMed:24695737, PubMed:24939902, PubMed:35585232). Auxiliary protein of DNA polymerase delta and epsilon, is involved in the control of DNA replication by increasing the polymerases' processivity during elongation of the leading strand (PubMed:35585232). Induces a robust stimulatory effect on the 3'-5' exonuclease and 3'- phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order to be able to stimulate APEX2. Plays a key role in DNA damage response (DDR) by being conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and DNA damage tolerance pathways (PubMed:24939902). Acts as a loading platform to recruit DDR proteins that allow completion of DNA replication after DNA damage and promote postreplication repair: monoubiquitinated PCNA leads to recruitment of translesion (TLS) polymerases, while 'Lys-63'-linked polyubiquitination of PCNA is involved in error-free pathway and employs recombination mechanisms to synthesize across the lesion (PubMed:24695737). |

| | |
|---|---|
| **Cellular Location** | Nucleus. Note=Colocalizes with CREBBP, EP300 and POLD1 to sites of DNA damage (PubMed:24939902). Forms nuclear foci representing sites of ongoing DNA replication and vary in morphology and number during S phase (PubMed:15543136). Co-localizes with SMARCA5/SNF2H and BAZ1B/WSTF at replication foci during S phase (PubMed:15543136). Together with APEX2, is redistributed in discrete nuclear foci in presence of oxidative DNA damaging agents |

## Images



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human lung squamous cell carcinoma tissue was performed with PCNA primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.