

CD105 Antibody

Mouse Monoclonal Antibody (Mab)

Catalog # AD80042

Product Information

| | |
|-------------------|------------------------|
| Application | IHC |
| Primary Accession | P17813 |
| Reactivity | Human |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG1 |
| Clone Names | 813A5D5 |
| Calculated MW | 70578 |

Additional Information

| | |
|-------------|--|
| Gene ID | 2022 |
| Gene Name | ENG |
| Other Names | Endoglin, CD105, ENG, END |
| Dilution | IHC~~Ready-to-use |
| Storage | Maintain refrigerated at 2-8°C. |
| Precautions | CD105 Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |


Protein Information

| | |
|----------|--|
| Name | ENG |
| Synonyms | END |
| Function | Vascular endothelium glycoprotein that plays an important role in the regulation of angiogenesis (PubMed: 21737454 , PubMed: 23300529). Required for normal structure and integrity of adult vasculature (PubMed: 7894484). Regulates the migration of vascular endothelial cells (PubMed: 17540773). Required for normal extraembryonic angiogenesis and for embryonic heart development (By similarity). May regulate endothelial cell shape changes in response to blood flow, which drive vascular remodeling and establishment of normal vascular morphology during angiogenesis (By similarity). May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors (PubMed: 1692830). Acts as a TGF-beta coreceptor and is involved in the TGF-beta/BMP signaling cascade that ultimately leads to the activation of SMAD transcription factors (PubMed: 21737454 , PubMed: 22347366 , PubMed: 23300529 , PubMed: 8370410). Required for GDF2/BMP9 signaling through SMAD1 in endothelial cells and modulates TGFβ1 signaling through SMAD3 (PubMed: 21737454 , PubMed: 22347366 , PubMed: 23300529). |

Cellular Location
Tissue Location

Cell membrane; Single-pass type I membrane protein
Detected on umbilical vein endothelial cells (PubMed:10625079). Detected in placenta (at protein level) (PubMed:1692830). Detected on endothelial cells (PubMed:1692830)

Images



Immunohistochemical analysis of paraffin-embedded colorectal carcinoma; tissue using AD80042 performed on the Abcarta® FAIP-48 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems (Abcepta: ADR005) was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.