

# **CD13**

Rabbit Monoclonal antibody(Mab)
Catalog # AD80057

#### **Product Information**

Application IHC-P
Primary Accession P15144
Reactivity Human
Host Rabbit
Clonality Monoclonal
Clone Names 548B9E9
Calculated MW 109540

### **Additional Information**

Gene ID 290 Gene Name ANPEP

Other Names Aminopeptidase N, AP-N, hAPN, 3.4.11.2, Alanyl aminopeptidase,

Aminopeptidase M, AP-M, Microsomal aminopeptidase, Myeloid plasma membrane glycoprotein CD13, gp150, CD13, ANPEP, APN, CD13, PEPN

**Dilution** IHC-P~~Ready-to-use

**Storage** Maintain refrigerated at 2-8°C.

**Precautions** CD13 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

### **Protein Information**

Name ANPEP

**Synonyms** APN, CD13, PEPN

**Function** Broad specificity aminopeptidase which plays a role in the final digestion of

peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Also involved in the processing of various peptides including peptide hormones, such as angiotensin III and IV, neuropeptides, and chemokines. May also be involved the cleavage of peptides bound to major histocompatibility complex class II molecules of antigen presenting cells. May have a role in angiogenesis and promote cholesterol crystallization. May have a role in amino acid transport by acting as binding partner of amino acid

transporter SLC6A19 and regulating its activity (By similarity).


**Cellular Location** Cell membrane; Single-pass type II membrane protein. Note=Also found as a

soluble form

**Tissue Location** Expressed in epithelial cells of the kidney, intestine, and respiratory tract;

granulocytes, monocytes, fibroblasts, endothelial cells, cerebral pericytes at the blood-brain barrier, synaptic membranes of cells in the CNS. Also expressed in endometrial stromal cells, but not in the endometrial glandular cells. Found in the vasculature of tissues that undergo angiogenesis and in malignant gliomas and lymph node metastases from multiple tumor types but not in blood vessels of normal tissues. A soluble form has been found in plasma. It is found to be elevated in plasma and effusions of cancer patients.

# **Images**



扁桃体

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.