

# ALK

Mouse Monoclonal antibody(Mab)

Catalog # AD80132

## Product Information

Application	IHC-P
Primary Accession	<a href="#">Q9UM73</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	137E9E8
Calculated MW	176442

## Additional Information

Gene ID	238
Gene Name	ALK
Other Names	ALK tyrosine kinase receptor, 2.7.10.1, Anaplastic lymphoma kinase, CD246, ALK {ECO:0000303   PubMed:9174053, ECO:0000312   HGNC:HGNC:427}
Dilution	IHC-P~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.
Precautions	ALK Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	ALK {ECO:0000303   PubMed:9174053, ECO:0000312   HGNC:HGNC:427}
Function	Neuronal receptor tyrosine kinase that is essentially and transiently expressed in specific regions of the central and peripheral nervous systems and plays an important role in the genesis and differentiation of the nervous system (PubMed: <a href="#">11121404</a> , PubMed: <a href="#">11387242</a> , PubMed: <a href="#">16317043</a> , PubMed: <a href="#">17274988</a> , PubMed: <a href="#">30061385</a> , PubMed: <a href="#">34646012</a> , PubMed: <a href="#">34819673</a> ). Also acts as a key thinness protein involved in the resistance to weight gain: in hypothalamic neurons, controls energy expenditure acting as a negative regulator of white adipose tissue lipolysis and sympathetic tone to fine-tune energy homeostasis (By similarity). Following activation by ALKAL2 ligand at the cell surface, transduces an extracellular signal into an intracellular response (PubMed: <a href="#">30061385</a> , PubMed: <a href="#">33411331</a> , PubMed: <a href="#">34646012</a> , PubMed: <a href="#">34819673</a> ). In contrast, ALKAL1 is not a potent physiological ligand for ALK (PubMed: <a href="#">34646012</a> ). Ligand-binding to the extracellular domain induces tyrosine kinase activation, leading to activation of the mitogen-activated protein kinase (MAPK) pathway (PubMed: <a href="#">34819673</a> ). Phosphorylates almost exclusively at the first tyrosine of

the Y-x-x-x-Y-Y motif (PubMed:[15226403](#), PubMed:[16878150](#)). Induces tyrosine phosphorylation of CBL, FRS2, IRS1 and SHC1, as well as of the MAP kinases MAPK1/ERK2 and MAPK3/ERK1 (PubMed:[15226403](#), PubMed:[16878150](#)). ALK activation may also be regulated by pleiotrophin (PTN) and midkine (MDK) (PubMed:[11278720](#), PubMed:[11809760](#), PubMed:[12107166](#), PubMed:[12122009](#)). PTN-binding induces MAPK pathway activation, which is important for the anti-apoptotic signaling of PTN and regulation of cell proliferation (PubMed:[11278720](#), PubMed:[11809760](#), PubMed:[12107166](#)). MDK-binding induces phosphorylation of the ALK target insulin receptor substrate (IRS1), activates mitogen-activated protein kinases (MAPKs) and PI3-kinase, resulting also in cell proliferation induction (PubMed:[12122009](#)). Drives NF-kappa-B activation, probably through IRS1 and the activation of the AKT serine/threonine kinase (PubMed:[15226403](#), PubMed:[16878150](#)). Recruitment of IRS1 to activated ALK and the activation of NF-kappa-B are essential for the autocrine growth and survival signaling of MDK (PubMed:[15226403](#), PubMed:[16878150](#)).

#### Cellular Location


Cell membrane; Single-pass type I membrane protein Note=Membrane attachment is essential for promotion of neuron-like differentiation and cell proliferation arrest through specific activation of the MAP kinase pathway.

#### Tissue Location

Expressed in brain and CNS. Also expressed in the small intestine and testis, but not in normal lymphoid cells

## Images

---



肺腺癌

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.