10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CD30

Mouse Monoclonal antibody(Mab)
Catalog # AD80137

Product Information

Application	IHC-P
Primary Accession	P28908
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	442F7G3
Calculated MW	63747

Additional Information


Gene ID	943
Other Names	Tumor necrosis factor receptor superfamily member 8 {ECO:0000312|HGNC:HGNC:11923}, CD30L receptor, Ki-1 antigen, Lymphocyte activation antigen CD30, CD30, TNFRSF8 (HGNC:11923)
Dilution	IHC-P~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.


Protein Information

Name	TNFRSF8 (HGNC:11923)
Function	Receptor for TNFSF8/CD30L (PubMed: 8391931). May play a role in the regulation of cellular growth and transformation of activated lymphoblasts. Regulates gene expression through activation of NF-kappa- B (PubMed:8999898).
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein
Tissue Location	[Isoform 2]: Detected in alveolar macrophages (at protein level).


Images

Immunohistochemical analysis of formalin-fixed and paraffin-embedded human appendix tissue was performed with CD30 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.





Immunohistochemical analysis of formalin-fixed and paraffin-embedded human appendix tissue was performed with CD30 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human hodgkin Lymphoma tissue was performed with CD30 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.