

PU.1

Mouse Monoclonal antibody(Mab)

Catalog # AD80158

Product Information

Application	IHC-P
Primary Accession	P17947
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	199E3D7
Calculated MW	31083

Additional Information

Gene ID	6688
Gene Name	SPI1
Other Names	Transcription factor PU.1, 31 kDa-transforming protein, SPI1
Dilution	IHC-P~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.
Precautions	PU.1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	SPI1
Function	<p>Pioneer transcription factor, which controls hematopoietic cell fate by decompacting stem cell heterochromatin and allowing other transcription factors to enter otherwise inaccessible genomic sites. Once in open chromatin, can directly control gene expression by binding genetic regulatory elements and can also more broadly influence transcription by recruiting transcription factors, such as interferon regulatory factors (IRFs), to otherwise inaccessible genomic regions (PubMed:23658224, PubMed:33951726). Transcriptionally activates genes important for myeloid and lymphoid lineages, such as CSF1R (By similarity). Transcriptional activation from certain promoters, possibly containing low affinity binding sites, is achieved cooperatively with other transcription factors. FCER1A transactivation is achieved in cooperation with GATA1 (By similarity). May be particularly important for the pro- to pre-B cell transition (PubMed:33951726). Binds (via the ETS domain) onto the purine-rich DNA core sequence 5'-GAGGAA-3', also known as the PU-box (PubMed:33951726). In vitro can bind RNA and interfere with pre-mRNA splicing (By similarity).</p>
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00237,

Tissue Location

ECO:0000269 | PubMed:33951726}

In the bone marrow, concentrated in hematopoietic stem cell, lymphoid progenitor, myeloid lineage (granulocyte macrophage progenitors, classical dendritic cells, monocytes) and B-cell clusters Among B-cells, predominantly expressed in pre-B1 cells (PubMed:33951726). Expressed in germinal center B-cells (PubMed:23166356).

Images



扁桃体

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.