

## EZH2

Rabbit Monoclonal antibody(Mab)

Catalog # AD80174

### Product Information

---

Application	IHC-P
Primary Accession	<a href="#">Q15910</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Clone Names	658B1F3
Calculated MW	85363

### Additional Information

---

Gene ID	2146
Gene Name	EZH2 ( <a href="#">HGNC:3527</a> )
Other Names	Histone-lysine N-methyltransferase EZH2, 2.1.1.356, ENX-1, Enhancer of zeste homolog 2, Lysine N-methyltransferase 6, EZH2 ( <a href="#">HGNC:3527</a> ), KMT6
Dilution	IHC-P~~N/A
Storage	Maintain refrigerated at 2-8°C.
Precautions	EZH2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

---

Name	EZH2 ( <a href="#">HGNC:3527</a> )
Synonyms	KMT6
Function	<p>Polycomb group (PcG) protein. Catalytic subunit of the PRC2/EED-EZH2 complex, which methylates 'Lys-9' (H3K9me) and 'Lys-27' (H3K27me) of histone H3, leading to transcriptional repression of the affected target gene. Able to mono-, di- and trimethylate 'Lys-27' of histone H3 to form H3K27me1, H3K27me2 and H3K27me3, respectively. Displays a preference for substrates with less methylation, loses activity when progressively more methyl groups are incorporated into H3K27, H3K27me0 &gt; H3K27me1 &gt; H3K27me2 (PubMed:<a href="#">22323599</a>, PubMed:<a href="#">30923826</a>). Compared to EZH1-containing complexes, it is more abundant in embryonic stem cells and plays a major role in forming H3K27me3, which is required for embryonic stem cell identity and proper differentiation. The PRC2/EED-EZH2 complex may also serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic repression systems. Genes repressed by the PRC2/EED-EZH2 complex include HOXC8, HOXA9, MYT1, CDKN2A and retinoic acid target genes. EZH2 can also methylate non-histone proteins such as the</p>

transcription factor GATA4 and the nuclear receptor RORA. Regulates the circadian clock via histone methylation at the promoter of the circadian genes. Essential for the CRY1/2-mediated repression of the transcriptional activation of PER1/2 by the CLOCK-BMAL1 heterodimer; involved in the di and trimethylation of 'Lys-27' of histone H3 on PER1/2 promoters which is necessary for the CRY1/2 proteins to inhibit transcription.

**Cellular Location**

Nucleus. Note=Localizes to the inactive X chromosome in trophoblast stem cells. {ECO:0000250|UniProtKB:Q61188}

**Tissue Location**

In the ovary, expressed in primordial follicles and oocytes and also in external follicle cells (at protein level) (PubMed:31451685). Expressed in many tissues (PubMed:14532106) Overexpressed in numerous tumor types including carcinomas of the breast, colon, larynx, lymphoma and testis (PubMed:14532106)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.