

IgG4

Rabbit Monoclonal antibody(Mab)

Catalog # AD80197

Product Information

Application	IHC-P
Primary Accession	P01861
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Clone Names	863G1E2
Calculated MW	43832


Additional Information

Gene Name	IGHG4 {ECO:0000303 PubMed:11340299, ECO:0000303 Ref.5}
Other Names	Immunoglobulin heavy constant gamma 4 {ECO:0000303 PubMed:11340299, ECO:0000303 Ref.6}, Ig gamma-4 chain C region, IGHG4 {ECO:0000303 PubMed:11340299, ECO:0000303 Ref.6}
Dilution	IHC-P~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.
Precautions	IgG4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	IGHG4 {ECO:0000303 PubMed:11340299, ECO:0000303 Ref.6}
Function	Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins- secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed: 20176268 , PubMed: 22158414). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed: 17576170 , PubMed: 20176268). [Isoform 1]: Secreted
Cellular Location	

Images



阑尾

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.