

# CD7 Antibody

Mouse Monoclonal Antibody (Mab) Catalog # AD80239

### **Product Information**

**Application** IHC **Primary Accession** P09564 Reactivity Human Host Mouse Clonality Monoclonal Isotype IgG1 **Clone Names** 634A6D3 Calculated MW 25409

## **Additional Information**

Gene ID 924 Gene Name CD7

Other Names T-cell antigen CD7, GP40, T-cell leukemia antigen, T-cell surface antigen Leu-9,

TP41, CD7, CD7

**Dilution** IHC~~Ready-to-use

**Storage** Maintain refrigerated at 2-8°C.

**Precautions** CD7 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

# **Protein Information**

Name CD7

**Function** Transmembrane glycoprotein expressed by T-cells and natural killer (NK)

cells and their precursors (PubMed:<u>7506726</u>). Plays a costimulatory role in T-cell activation upon binding to its ligand K12/SECTM1 (PubMed:<u>10652336</u>). In turn, mediates the production of cytokines such as IL-2 (PubMed:<u>1709867</u>).

On resting NK-cells, CD7 activation results in a significant induction of  $% \left\{ 1\right\} =\left\{ 1\right\} =$ 


interferon-gamma levels (PubMed: 7506726).

**Cellular Location** Membrane; Single-pass type I membrane protein.

**Tissue Location** Expressed on T-cells and natural killer (NK) cells and their precursors.

#### **Images**

Immunohistochemical analysis of paraffin-embedded human tonsil tissue using AD80239 performed on the Abcarta® FAIP-48 Fully automated IHC platform. Tissue



was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6. 0). Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSeeTM Detection Systems(Abcepta: ADR005) was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.