

## Oligo-2 Antibody

Mouse Monoclonal Antibody (Mab) Catalog # AD80268

## **Product Information**

Application IHC
Primary Accession Q13516
Reactivity Human
Host Mouse
Clonality Monoclonal
Clone Names 294A3D1
Calculated MW 32385

## **Additional Information**

**Gene ID** 10215 **Gene Name** OLIG2

Other Names Oligodendrocyte transcription factor 2, Oligo2, Class B basic helix-loop-helix

protein 1, bHLHb1, Class E basic helix-loop-helix protein 19, bHLHe19, Protein kinase C-binding protein 2, Protein kinase C-binding protein RACK17, OLIG2,

BHLHB1, BHLHE19, PRKCBP2, RACK17

**Dilution** IHC~~1:100~500

**Storage** Maintain refrigerated at 2-8°C.

**Precautions** Oligo-2 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

## **Protein Information**

Name OLIG2

**Synonyms** BHLHB1, BHLHE19, PRKCBP2, RACK17

**Function** Required for oligodendrocyte and motor neuron specification in the spinal

cord, as well as for the development of somatic motor neurons in the hindbrain. Functions together with ZNF488 to promote oligodendrocyte differentiation. Cooperates with OLIG1 to establish the pMN domain of the embryonic neural tube. Antagonist of V2 interneuron and of NKX2-2-induced

V3 interneuron development.

Cellular Location Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00981}. Cytoplasm. Note=The

NLS contained in the bHLH domain could be masked in the native form and translocation to the nucleus could be mediated by interaction either with

class E bHLH partner protein or with NKX2-2.

**Tissue Location** Expressed in the brain, in oligodendrocytes. Strongly expressed in

oligodendrogliomas, while expression is weak to moderate in astrocytomas.

Expression in glioblastomas highly variable

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.