10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

C1q

Mouse Monoclonal antibody(Mab)
Catalog # AD80317

Product Information

Application	IHC-P
Primary Accession	Q9NPY3
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	652A2F4
Calculated MW	68560

Additional Information


Gene ID	22918
Gene Name	CD93
Other Names	Complement component C1q receptor, C1q/MBL/SPA receptor, C1qR, C1qR(p), C1qRp, CDw93, Complement component 1 q subcomponent receptor 1, Matrix-remodeling-associated protein 4, CD93, CD93, C1QR1, MXRA4
Dilution	IHC-P~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.
Precautions	C1q Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CD93
Synonyms	C1QR1, MXRA4
Function	Cell surface receptor that plays a role in various physiological processes including inflammation, phagocytosis, and cell adhesion. Plays a role in phagocytosis and enhances the uptake of apoptotic cells and immune complexes by acting as a receptor for defense collagens including surfactant protein A/SFTPA1, C1q, and mannose-binding lectin (MBL2) (PubMed:7977768). Plays a role in the regulation of endothelial cell function and adhesion by activating angiogenesis (PubMed:24809468). Mechanistically, exerts its angiogenic function by associating with beta-dystroglycan, leading to SRC- dependent phosphorylation and subsequent recruitment of CBL. In turn, CBL provides a docking site for downstream signaling components, such as CRKL to enhance cell migration (PubMed:26848865). Participates in angiogenesis also by acting as a receptor for the ECM pan-endothelial glycoprotein multimerin-2/MMRN2 and IGFBP7 ligands (PubMed:28671670, PubMed:36265539, PubMed:38218180). Both ligands play a non-redundant

	role in CD93-mediated endothelial cell function (PubMed:38218180). Acts as a key regulator of endothelial barrier function through modulating VEGFR2 function (By similarity).
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Highly expressed in endothelial cells, platelets, cells of myeloid origin, such as monocytes and neutrophils. Not expressed in cells of lymphoid origin

Images



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human appendix tissue was performed with C1q primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.