10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CK8&18

Rabbit Monoclonal antibody(Mab)
Catalog # AD80355

Product Information

Application	IHC-P
Primary Accession	P05787
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Clone Names	459G1A4
Calculated MW	53704


Additional Information

Gene ID	3856
Gene Name	KRT8
Other Names	Keratin, type II cytoskeletal 8, Cytokeratin-8, CK-8, Keratin-8, K8, Type-II keratin Kb8, KRT8, CYK8
Dilution	IHC-P~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.
Precautions	CK8&18 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	KRT8 (HGNC:6446)
Synonyms	CYK8
Function	Structural component of intermediate filaments in simple epithelial cells. Together with its obligate type I partner KRT18, contributes to the formation of the keratin intermediate filament network that provides mechanical stability and resilience to epithelial cells (PubMed:26823019). Also involved in maintaining epithelial integrity and protecting cells from non-mechanical stress, such as apoptosis (By similarity). Also required for ARHGEF40-mediated actin stress fiber formation and tensional force-induced stress fiber formation and reinforcement (PubMed:26823019). In striated muscle, together with KRT19, may contribute to linking the contractile apparatus to dystrophin at the costameres (PubMed:16000376).
Cellular Location	Cytoplasm. Nucleus, nucleoplasm {ECO:0000250|UniProtKB:Q10758}. Nucleus matrix {ECO:0000250|UniProtKB:Q10758}. Cytoplasm, cytoskeleton
Tissue Location	Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma membrane in structures that contain dystrophin and spectrin. Expressed in gingival mucosa and hard palate of the oral cavity.


Images




Immunohistochemical analysis of formalin-fixed and paraffin-embedded human breast carcinoma tissue was performed with CK8&18 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human ovarian carcinoma tissue was performed with CK8&18 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human appendix tissue was performed with CK8&18 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human prostatic adenocarcinoma tissue was performed with CK8&18 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.