10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Arginase-1

Rabbit Monoclonal antibody(Mab)
Catalog # AD80387

Product Information

Application	IHC-P
Primary Accession	P05089
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Clone Names	114G7B3
Calculated MW	34735

Additional Information


Gene ID	383
Gene Name	ARG1
Other Names	Arginase-1, 3.5.3.1, Liver-type arginase, Type I arginase, ARG1
Dilution	IHC-P~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.
Precautions	Arginase-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	ARG1
Function	Catalyzes the final step of the urea cycle by hydrolyzing L- arginine into urea, a nitrogenous waste product, and L-ornithine, a precursor to proline and polyamines; the urea cycle occurs primarily in the liver and, to a lesser extent, in the kidneys.
Cellular Location	Cytoplasm, cytosol. Cytoplasmic granule. Note=Localized in azurophil granules of neutrophils (PubMed:15546957)
Tissue Location	Within the immune system, initially reported to be selectively expressed in granulocytes (polymorphonuclear leukocytes [PMNs]) (PubMed:15546957). Also detected in macrophages in mycobacterial granulomas (PubMed:23749634). Expressed in group2 innate lymphoid cells (ILC2s) during lung disease (PubMed:27043409)

Images

Immunohistochemical analysis of formalin-fixed and paraffin-embedded human liver carcinoma tissue was



performed with Arginase-1 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human liver tissue was performed with Arginase-1 primary antibody, followed by peroxidase-conjugated secondary antibody and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.