

Granzyme B Antibody

Mouse Monoclonal Antibody (Mab)

Catalog # AD80412

Product Information

Application	IHC
Primary Accession	P10144
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG
Clone Names	974E3A1
Calculated MW	27716

Additional Information

Gene ID	3002
Gene Name	GZMB
Other Names	Granzyme B, 3.4.21.79, C11, CTLA-1, Cathepsin G-like 1, CTSG1, Cytotoxic T-lymphocyte proteinase 2, Lymphocyte protease, Fragmentin-2, Granzyme-2, Human lymphocyte protein, HLP, SECT, T-cell serine protease 1-3E, GZMB, CGL1, CSPB, CTLA1, GRB
Dilution	IHC~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.
Precautions	Granzyme B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	GZMB {ECO:0000303 PubMed:32188940, ECO:0000312 HGNC:HGNC:4709}
Function	Abundant protease in the cytosolic granules of cytotoxic T- cells and NK-cells which activates caspase-independent pyroptosis when delivered into the target cell through the immunological synapse (PubMed: 1985927 , PubMed: 3262682 , PubMed: 3263427). It cleaves after Asp (PubMed: 1985927 , PubMed: 8258716). Once delivered into the target cell, acts by catalyzing cleavage of gasdermin-E (GSDME), releasing the pore- forming moiety of GSDME, thereby triggering pyroptosis and target cell death (PubMed: 31953257 , PubMed: 32188940). Seems to be linked to an activation cascade of caspases (aspartate-specific cysteine proteases) responsible for apoptosis execution. Cleaves caspase-3, -9 and -10 (CASP3, CASP9 and CASP10, respectively) to give rise to active enzymes mediating apoptosis (PubMed: 9852092). Cleaves and activates CASP7 in response to bacterial infection, promoting plasma membrane repair (By similarity).

Cellular Location

Secreted. Cytolytic granule. Note=Delivered into the target cell by perforin (PubMed:20038786).

Images



Immunohistochemical analysis of paraffin-embedded human spleen tissue using AD80412 performed on the Abcarta® FAIP-48 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6. 0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems (Abcepta: ADR005) was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.