10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Melanoma(Pan)

Mouse Monoclonal antibody(Mab)
Catalog # AD80496

Product Information


Application	IHC-P
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	666F5C1

Additional Information

Gene ID	10215
Gene Name	OLIG2
Other Names	Oligodendrocyte transcription factor 2, Oligo2, Class B basic helix-loop-helix protein 1, bHLHb1, Class E basic helix-loop-helix protein 19, bHLHe19, Protein kinase C-binding protein 2, Protein kinase C-binding protein RACK17, OLIG2, BHLHB1, BHLHE19, PRKCBP2, RACK17
Dilution	IHC-P~~Ready-to-use
Storage	Maintain refrigerated at 2-8°C.
Precautions	Oligo-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Synonyms	BHLHB1, BHLHE19, PRKCBP2, RACK17
Function	Required for oligodendrocyte and motor neuron specification in the spinal cord, as well as for the development of somatic motor neurons in the hindbrain. Functions together with ZNF488 to promote oligodendrocyte differentiation. Cooperates with OLIG1 to establish the pMN domain of the embryonic neural tube. Antagonist of V2 interneuron and of NKX2-2-induced V3 interneuron development.
Cellular Location	Nucleus {ECO:0000255|PROSITE- ProRule:PRU00981}. Cytoplasm. Note=The NLS contained in the bHLH domain could be masked in the native form and translocation to the nucleus could be mediated by interaction either with class E bHLH partner protein or with NKX2-2
Tissue Location	Expressed in the brain, in oligodendrocytes. Strongly expressed in oligodendrogliomas, while expression is weak to moderate in astrocytomas. Expression in glioblastomas highly variable.

Images

Immunohistochemical analysis of formalin-fixed and paraffin-embedded human melanoma tissue was performed with Melanoma(Pan) primary antibody, followed by peroxidase-conjugated secondary antibody



and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.