

CD14 antibody

Mouse Monoclonal Antibody (Mab)

Catalog # AD80513

Product Information

Application	IHC
Primary Accession	P08571
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	607J7U4
Calculated MW	40076

Additional Information

Gene ID	929
Other Names	Monocyte differentiation antigen CD14, Myeloid cell-specific leucine-rich glycoprotein, CD14, Monocyte differentiation antigen CD14, urinary form, Monocyte differentiation antigen CD14, membrane-bound form, CD14
Dilution	IHC~~1:100~500
Storage	Maintain refrigerated at 2-8°C.

Protein Information

Name	CD14
Function	Coreceptor for bacterial lipopolysaccharide (PubMed: 1698311 , PubMed: 23264655). In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the LY96/TLR4 complex, thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed: 20133493 , PubMed: 22265692 , PubMed: 23264655). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed: 8612135). Acts as a coreceptor for TLR2:TLR6 heterodimer in response to diacylated lipopeptides and for TLR2:TLR1 heterodimer in response to triacylated lipopeptides, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway (PubMed: 16880211). Binds electronegative LDL (LDL(-)) and mediates the cytokine release induced by LDL(-) (PubMed: 23880187).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor. Secreted. Membrane raft. Golgi apparatus. Note=Secreted forms may arise by cleavage of the GPI anchor.
Tissue Location	Detected on macrophages (at protein level) (PubMed:1698311). Expressed strongly on the surface of monocytes and weakly on the surface of granulocytes; also expressed by most tissue macrophages.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.