10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



# HSP27

Mouse Monoclonal antibody(Mab)
Catalog # AD80518

#### **Product Information**

Application IHC-P
Primary Accession P04792
Reactivity Human
Host Mouse
Clonality Monoclonal
Clone Names 663K0B7
Calculated MW 22783

## **Additional Information**

Gene ID 3315

Other Names Heat shock protein beta-1, HspB1, 28 kDa heat shock protein,

Estrogen-regulated 24 kDa protein, Heat shock 27 kDa protein, HSP 27,

Stress-responsive protein 27, SRP27, HSPB1, HSP27, HSP28

**Dilution** IHC-P~~Ready-to-use

**Storage** Maintain refrigerated at 2-8°C.

### **Protein Information**

Name HSPB1

**Synonyms** HSP27, HSP28

**Function** Small heat shock protein which functions as a molecular chaperone

probably maintaining denatured proteins in a folding- competent state (PubMed:10383393, PubMed:20178975). Plays a role in stress resistance and actin organization (PubMed:19166925). Through its molecular chaperone activity may regulate numerous biological processes including the phosphorylation and the axonal transport of neurofilament proteins

(PubMed:23728742).

**Cellular Location** Cytoplasm. Nucleus Cytoplasm, cytoskeleton, spindle Note=Cytoplasmic in

interphase cells. Colocalizes with mitotic spindles in mitotic cells. Translocates to the nucleus during heat shock and resides in sub-nuclear structures known


as SC35 speckles or nuclear splicing speckles.

**Tissue Location** Detected in all tissues tested: skeletal muscle, heart, aorta, large intestine,

small intestine, stomach, esophagus, bladder, adrenal gland, thyroid, pancreas, testis, adipose tissue, kidney, liver, spleen, cerebral cortex, blood serum and cerebrospinal fluid. Highest levels are found in the heart and in

tissues composed of striated and smooth muscle.

# **Images**



乳腺癌

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.