

MLH1 antibody

Mouse Monoclonal Antibody (Mab)

Catalog # AD80539

Product Information

Application	IHC-P
Primary Accession	P40692
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Clone Names	451I3V0
Calculated MW	84601

Additional Information

Gene ID	4292
Other Names	DNA mismatch repair protein Mlh1, MutL protein homolog 1, MLH1, COCA2
Dilution	IHC-P~~N/A
Storage	Maintain refrigerated at 2-8°C.

Protein Information

Name	MLH1
Synonyms	COCA2
Function	<p>Heterodimerizes with PMS2 to form MutL alpha, a component of the post-replicative DNA mismatch repair system (MMR). DNA repair is initiated by MutS alpha (MSH2-MSH6) or MutS beta (MSH2-MSH3) binding to a dsDNA mismatch, then MutL alpha is recruited to the heteroduplex. Assembly of the MutL-MutS-heteroduplex ternary complex in presence of RFC and PCNA is sufficient to activate endonuclease activity of PMS2. It introduces single-strand breaks near the mismatch and thus generates new entry points for the exonuclease EXO1 to degrade the strand containing the mismatch. DNA methylation would prevent cleavage and therefore assure that only the newly mutated DNA strand is going to be corrected. MutL alpha (MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. Also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. Heterodimerizes with MLH3 to form MutL gamma which plays a role in meiosis.</p>
Cellular Location	Nucleus. Chromosome. Note=Recruited to chromatin in a MCM9- dependent manner.
Tissue Location	Colon, lymphocytes, breast, lung, spleen, testis, prostate, thyroid, gall bladder

and heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.