

CD208 antibody

Catalog # AD80576

Product Information

Application	IHC-P
Primary Accession	<u>Q9UQV4</u>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Clone Names	AB208
Calculated MW	44346

Additional Information

Gene ID	27074
Dilution	IHC-P~~N/A
Storage	Maintain refrigerated at 2-8°C.

Protein Information

Name	LAMP3
Synonyms Function	DCLAMP, TSC403 Lysosomal membrane glycoprotein which plays a role in the unfolded protein response (UPR) that contributes to protein degradation and cell survival during proteasomal dysfunction (PubMed: <u>25681212</u>). Plays a role in the process of fusion of the lysosome with the autophagosome, thereby modulating the autophagic process (PubMed: <u>24434718</u>). Promotes hepatocellular lipogenesis through activation of the PI3K/Akt pathway (PubMed: <u>29056532</u>). May also play a role in dendritic cell function and in adaptive immunity (PubMed: <u>9768752</u>).
Cellular Location	Cell surface. Lysosome membrane; Single-pass type I membrane protein. Cytoplasmic vesicle membrane; Single-pass type I membrane protein. Early endosome membrane; Single-pass type I membrane protein. Note=During dendritic cell maturation, detected on cytoplasmic vesicles (the MHC II compartment) that contain MHC II proteins, LAMP1, LAMP2 and LAMP3 (PubMed:9768752). Detected on lysosomes in mature dendritic cells (PubMed:9768752).
Tissue Location	Detected in tonsil interdigitating dendritic cells, in spleen, lymph node, Peyer's patches in the small instestine, in thymus medulla and in B-cells (at protein level). Expressed in lymphoid organs and dendritic cells. Expressed in lung. Up-regulated in carcinomas of the esophagus, colon, rectum, ureter, stomach, breast, fallopian tube, thyroid and parotid tissues

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.