

Folate Binding Protein(FBP)抗体试剂(免疫组织化学)

Rabbit Monoclonal antibody(Mab)
Catalog # AD80605

Product Information

Application IHC-P
Primary Accession P15328
Reactivity Human
Host Rabbit
Clonality Monoclonal
Clone Names 441I2C4
Calculated MW 29819

Additional Information

Gene ID 2348

Other Names Folate receptor alpha, FR-alpha, Adult folate-binding protein, FBP, Folate

receptor 1, Folate receptor, adult, KB cells FBP, Ovarian tumor-associated

antigen MOv18, FOLR1, FOLR

Dilution IHC-P~~N/A

Storage Maintain refrigerated at 2-8°C.

Protein Information

Name FOLR1

Synonyms FOLR

Function Binds to folate and reduced folic acid derivatives and mediates delivery of

5-methyltetrahydrofolate and folate analogs into the interior of cells

(PubMed: 19074442, PubMed: 23851396, PubMed: 23934049,

PubMed: <u>2527252</u>, PubMed: <u>8033114</u>, PubMed: <u>8567728</u>). Has high affinity for

folate and folic acid analogs at neutral pH (PubMed:23851396,

PubMed:23934049, PubMed:2527252, PubMed:8033114, PubMed:8567728).

Exposure to slightly acidic pH after receptor endocytosis triggers a conformation change that strongly reduces its affinity for folates and mediates their release (PubMed:8567728). Required for normal embryonic

development and normal cell proliferation (By similarity).

Cellular Location Cell membrane; Lipid-anchor, GPI-anchor Apical cell membrane; Lipid-anchor,

GPI- anchor Basolateral cell membrane; Lipid-anchor, GPI-like-anchor. Secreted Cytoplasmic vesicle. Cytoplasmic vesicle, clathrin-coated vesicle. Endosome. Note=Endocytosed into cytoplasmic vesicles and then recycled to

the cell membrane

Tissue Location Primarily expressed in tissues of epithelial origin. Expression is increased in

malignant tissues. Expressed in kidney, lung and cerebellum. Detected in

placenta and thymus epithelium.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.